

Original Investigation

Vegetarian Diets and Blood Pressure

A Meta-analysis

Yoko Yokoyama, PhD, MPH; Kunihiro Nishimura, MD, PhD, MPH; Neal D. Barnard, MD; Misa Takegami, RN, PhD, MPH; Makoto Watanabe, MD, PhD; Akira Sekikawa, MD, PhD; Tomonori Okamura, MD, PhD; Yoshihiro Miyamoto, MD, PhD

IMPORTANCE Previous studies have suggested an association between vegetarian diets and lower blood pressure (BP), but the relationship is not well established.

OBJECTIVE To conduct a systematic review and meta-analysis of controlled clinical trials and observational studies that have examined the association between vegetarian diets and BP.

DATA SOURCES MEDLINE and Web of Science were searched for articles published in English from 1946 to October 2013 and from 1900 to November 2013, respectively.

STUDY SELECTION All studies met the inclusion criteria of the use of (1) participants older than 20 years, (2) vegetarian diets as an exposure or intervention, (3) mean difference in BP as an outcome, and (4) a controlled trial or observational study design. In addition, none met the exclusion criteria of (1) use of twin participants, (2) use of multiple interventions, (3) reporting only categorical BP data, or (4) reliance on case series or case reports.


DATA EXTRACTION AND SYNTHESIS Data collected included study design, baseline characteristics of the study population, dietary data, and outcomes. The data were pooled using a random-effects model.

MAIN OUTCOMES AND MEASURES Net differences in systolic and diastolic BP associated with the consumption of vegetarian diets were assessed.

RESULTS Of the 258 studies identified, 7 clinical trials and 32 observational studies met the inclusion criteria. In the 7 controlled trials (a total of 311 participants; mean age, 44.5 years), consumption of vegetarian diets was associated with a reduction in mean systolic BP (−4.8 mm Hg; 95% CI, −6.6 to −3.1; $P < .001$; $I^2 = 0$; $P = .45$ for heterogeneity) and diastolic BP (−2.2 mm Hg; 95% CI, −3.5 to −1.0; $P < .001$; $I^2 = 0$; $P = .43$ for heterogeneity) compared with the consumption of omnivorous diets. In the 32 observational studies (a total of 21 604 participants; mean age, 46.6 years), consumption of vegetarian diets was associated with lower mean systolic BP (−6.9 mm Hg; 95% CI, −9.1 to −4.7; $P < .001$; $I^2 = 91.4$; $P < .001$ for heterogeneity) and diastolic BP (−4.7 mm Hg; 95% CI, −6.3 to −3.1; $P < .001$; $I^2 = 92.6$; $P < .001$ for heterogeneity) compared with the consumption of omnivorous diets.

CONCLUSIONS AND RELEVANCE Consumption of vegetarian diets is associated with lower BP. Such diets could be a useful nonpharmacologic means for reducing BP.

JAMA Intern Med. 2014;174(4):577-587. doi:10.1001/jamainternmed.2013.14547
Published online February 24, 2014.

 Supplemental content at
jamainternalmedicine.com

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Yoko Yokoyama, PhD, MPH, Department of Preventive Medicine and Epidemiologic Informatics, National Cerebral and Cardiovascular Center, 5-7-1 Fujishirodai, Suita-city, Osaka 565-8565, Japan (yyokoyama-kyt@umin.ac.jp).

The relationship between blood pressure (BP) and cardiovascular disease risk is continuous, consistent, and independent of other risk factors.¹ According to Lewington et al,² in individuals aged 40 to 70 years, each increment of 20 mm Hg in systolic BP or 10 mm Hg in diastolic BP is associated with more than twice the risk of cardiovascular disease across the BP range from 115/75 to 185/115 mm Hg.

A substantial body of evidence supports the role of modifiable factors, including diet, body weight, physical activity, and alcohol intake, in the risk of developing hypertension.³ Dietary modifications have been shown³ to be particularly effective in preventing and managing hypertension.

Vegetarian diets are defined as dietary patterns that exclude or rarely include meats; some vegetarian diets include dairy products, eggs, and fish. All vegetarian diets emphasize foods of plant origin, particularly vegetables, grains, legumes, and fruits. In observational studies,^{4,5} consumption of vegetarian diets is associated with a lower prevalence of hypertension. Although some randomized clinical trials^{6,7} have found that adoption of a vegetarian diet reduces BP, others^{8,9} have not yielded similar results. To our knowledge, the available evidence regarding the association between vegetarian diets and BP has not been subjected to meta-analysis. To clarify

the nature of this association and provide a valid estimate of the effect size regarding the effects of consumption of vegetarian diets on BP, both of which could prove useful in formulating dietary guidance, we performed a meta-analysis of studies that had examined associations between vegetarian diets and BP.

Methods

Data Sources and Search Strategy

The electronic search strategy is shown in the Supplement (eTable 1). MEDLINE and Web of Science were searched for articles published in English from January 1, 1946, to November 7, 2013, and from January 1, 1900, to November 7, 2013, respectively, containing 1 or more of the keywords or phrases for vegetarian diets (*plant-based diet* or *diet*, *vegetarian* or *vegetarian diets* or *vegetarianism* or *diets*, *vegan* or *vegan diets*) and for blood pressure (*blood pressure* or *hypertension*). The reference lists of the retrieved articles were subsequently reviewed for identification of additional articles. If necessary, the relevant authors were contacted by the investigators to acquire missing information (Figure 1).


Study Selection

Two reviewers (Y.Y. and M.T.) independently scanned the retrieved abstracts to identify studies that met the following inclusion criteria: (1) use of a sample of participants older than 20 years; (2) an intervention or exposure consisting of a vegetarian diet, defined as a diet generally excluding or rarely including meats; these may include *semivegetarian diets*, defined as rarely including meat; *vegan diets*, defined as omitting all animal products; or *vegetarian diets* that include some animal products as indicated by the terms *lacto* (dairy products), *ovo* (eggs), or *pesco* (fish); (3) collection of sufficient data to allow calculation of mean differences in systolic/diastolic BP between individuals consuming a vegetarian diet and those consuming a referent or control diet; and (4) use of a controlled trial or observational study design. The exclusion criteria were (1) use of a sample consisting of twins; (2) use of multiple interventions (ie, use of lifestyle interventions in addition to dietary interventions); (3) reporting only categorical BP data; or (4) reliance on case series or case reports.

Data Extraction and Quality Assessment

For each study, data regarding systolic and diastolic BP and variance measures; study methodology and sample size; baseline characteristics of the study population, including mean age, sex (proportion of men), BP, antihypertensive medication use, body mass index (BMI) (calculated as weight in kilograms divided by height in meters squared), alcohol intake, and dietary data (including type of diets examined and duration of their consumption); and outcomes, including adjustment factors used for each analytic model, BP measurements, and dietary measurements, were extracted. Mean values for baseline age, proportion of men, systolic and diastolic BP, BMI, and alcohol intake were calculated.

Figure 1. Study Flow Diagram



Selection of clinical trials and observational studies for meta-analysis of association between vegetarian diets and blood pressure.

Table 1. Designs and Population Characteristics of Clinical Trials of Vegetarian Diets and BP

Source	Country	Study Design (Duration)	No. of Participants	Mean Age, y	Male Sex, %	Mean BP, mm Hg		Using Medication, %	Mean BMI	Alcohol Intake	Intervention, Food Preparation ^a
						Systolic	Diastolic				
Ferdowsian et al, ¹² 2010	US	P, O (22 wk)	113	44.4	17.7	117.8	79.7	NR	NR	Participants with active alcohol abuse excluded	Vegan, yes (2/d)
Nicholson et al, ¹³ 1999	US	P, O (12 wk) ^b	11	54.3	54.5	141.3	84.7	81.8	NR	Individuals using alcohol regularly excluded	Vegan, yes
Sciarrone et al, ¹⁴ 1993	Australia	P, O (6 wk) ^b	20	41.0	100	134.2	77.2	None	25.3	Individuals using > 20 g of ethanol/d excluded	Lacto-ovo, yes
Hakala and Karvetti, ¹⁵ 1989	Finland	P, O (52 wk) ^b	73	38.0	24.7	129.9	85.0	None	34.4	Veg/Cont, 2%/2% energy	Lacto, no
Kestin et al, ⁸ 1989	Australia	C, O (6 wk) ^b	17	44.0	100	128.0	79.0	None	25.5	Veg/Cont, 4.2%/4.8% energy	Lacto-ovo, yes (major sources of protein and fat)
Margetts et al, ⁷ 1986	Australia	C, O (6 wk) ^b	39	49.9	71.8	155.4	99.9	None	27.6	Participants asked to not alter alcohol consumption	Lacto-ovo, yes (meat substitutes)
Rouse et al, ⁶ 1983	Australia	C, O (6 wk) ^b	38	40.1	50.0	127.7	76.4	None	23.7	Participants asked to not alter alcohol consumption	Lacto-ovo, yes (2/d)

Abbreviations: BP, blood pressure; BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); C, crossover; Cont, control; NR, not reported; O, open label; P, parallel; US, United States; Veg, vegetarian.

^a Vegan diets, defined as omitting all animal products, or vegetarian diets that

include some animal products as indicated by the terms *lacto* (dairy products) and *ovo* (eggs).

^b Randomized clinical trial.

Data Synthesis and Analysis

Mean differences in systolic and diastolic BP between groups consuming vegetarian or comparison diets were calculated. The pooled SE for the net difference in BP associated with the consumption of a vegetarian diet was obtained or, when not given, estimated using the method of Follmann et al,¹⁰ assuming a correlation of 0.50 between the baseline and final BP values (parallel design) or between the BP values during the intervention and control periods (crossover design). For studies comparing more than 1 exposure group or treatment arm, such as those comparing vegan and lacto-ovo-vegetarian groups, a pooled effect was calculated for each study using a random-effects model and then used to conduct the overall calculation.

Estimates of net change in BP associated with the consumption of vegetarian diets were combined using a random-effects model, which assigns a weight to each study on the basis of an individual study's inverse variance. Overall estimates were derived for controlled trials and observational studies separately, using the study as the unit of analysis. Estimates of BP differences were reported within 95% CIs. Differences were considered significant at 2-sided $P < .05$.

Stratified analyses by mean age, sex, BMI, diet type, sample size, duration of vegetarian diet consumption, antihypertensive medication use, baseline hypertensive status, and location (country) were performed separately for controlled trials and observational studies. As a sensitivity analysis, we conducted a one-study-removed analysis to assess the effect of each study on the combined effect. Calculation of I^2 and meta-regression was performed with subgroups, using the study as the unit of analysis to assess heterogeneity among studies.¹¹

Funnel plots were developed and examined to identify publication bias, and the Egger test was performed to assess

the relationship between sample size and effect size. The trim-and-fill method was used to adjust for publication bias. The trim-and-fill method determines where missing studies are likely to fall, adds them to the analysis, and then recomputes the combined effect. These analyses were conducted separately for controlled trials and observational studies. All analyses were performed using Comprehensive Meta-analysis, version 2, software (Biostat).

Results

Search Results

The search of the MEDLINE and Web of Science databases led to the retrieval of 258 studies. Of these, 7 clinical trials and 32 observational studies met the inclusion criteria (Figure 1).

Study Characteristics and Quality

Clinical Trials

Seven clinical trials were identified (Table 1).^{6-8,12-15} The 7 trials included a total of 311 participants (median sample size, 38; range, 11-113), with a mean age of 44.5 years (range, 38.0-54.3 years). All were open (nonmasked) controlled trials conducted for 6 or more weeks (mean, 15.7 weeks). Of these, 6 were randomized clinical trials.^{6-8,13-15} As shown in Table 1, several participants in 1 clinical trial¹³ used antihypertensive medication. All except 1 study¹⁵ provided foods to the participants. Vegan diets were examined in 2 trials,^{12,13} a lacto-vegetarian diet in 1,¹⁵ and lacto-ovo-vegetarian diets in 4.^{6-8,14} Four studies¹²⁻¹⁵ used parallel designs, and 3 trials⁶⁻⁸ used crossover designs. All studies^{6-8,12-15} reported repeated BP measurements. Adjustments for potential confounders for each trial are shown in the Supplement (eTable 2).

Table 2. Designs and Population Characteristics of Observational Cross-Sectional Studies of Vegetarian Diets and BP

Source	Country	No. of Participants	Mean Age, y	Male Sex, %	Mean BP, mm Hg		Using Antihypertensive Medication, %	Mean BMI	Alcohol Intake (Veg/Cont)	Exposure, Duration of Exposure ^a
					Systolic	Diastolic				
Kim and Bae, ¹⁶ 2012	South Korea	107	62.6	0	141.4	85.3	NR	23.8	Mild alcohol drinker, 0%/20.8% (overall, 9.4%)	Lacto-ovo, >20 y
Pettersen et al, ¹⁷ 2012	US, Canada	431	62.8	36.7	125.1	74.8	24.8	26.7	Minimal or absent in both groups, 6.75%	Mixed (vegan, lacto-ovo), >1 y
Yang et al, ¹⁸ 2012	China	295	33.3	100	120.5	74.7	None	24.0	Regular drinking excluded	Lacto, >5 y
Chen et al, ¹⁹ 2011	Taiwan	363	51.9	0	121.4	71.6	None	23.1	NR	Lacto-ovo, >1 y
Fernandes Dourado et al, ²⁰ 2011	Brazil	87	40.0	58.6	120.8	75.9	NR	24.3	0/12.1%	Lacto-ovo, >1 y (mean, 16 y)
Rodenas et al, ²¹ 2011	Spain	26	68.4	0	135.4	73.4	None	24.2	NR	Mixed (meatless), current
Yang et al, ²² 2011	China	300	33.3	100	120.3	74.4	None	23.9	Excluded	Lacto-ovo, >1 y (mean, 10.4 y)
Lin et al, ²³ 2010	Taiwan	204	46.1	0	117.3	75.2	NR	23.4	NR	Vegan, mean, 17.8 y
Pitla and Nagalla, ²⁴ 2009	India	Male, 29; female, 23	46.2; 45.4	100; 0	125.0; 117.8	81.7; 76.1	NR	24.8; 24.9	17%; 0%	Lacto, lifetime
Nakamoto et al, ²⁵ 2008	Japan	Male, 49; female, 73	44.1; 45.4	100; 0	124.9; 113.4	78.6; 68.4	NR	22.5; 21.4	NR	Mixed (lacto-ovo, semi), current
Slavíček et al, ²⁶ 2008	Czech	396	47.4	34.1	125.5	75.8	NR	23.6	NR	Lacto-ovo, >5 y
Fontana et al, ²⁷ 2007	US	42	53.1	61.9	118.0	70.5	None	23.9	NR	Vegan, >2 y (mean, 4.4 y)
Teixeira et al, ²⁸ 2007	Brazil	201	47.0	47.8	122.0	81.0	11.4	25.3	Veg < Cont	Mixed (vegan, lacto, lacto-ovo, pescos), >5 y (mean, 19 y)
Sebeková et al, ²⁹ 2006	Slovakia	136	37.5	36.0	112.3	70.9	None	23.1	NR	Lacto-ovo, >2 y (mean, 10.3 y)
Su et al, ³⁰ 2006	Taiwan	118	58.4	0	127.2	72.0	None	23.3	Regular drinking excluded	Mixed (vegan, lacto), >5 y (mean, 10.4 y)
Goff et al, ³¹ 2005	UK	46	35.5	46.7	118.7	68.6	NR	23.1	NR	Vegan, >3 y
Appleby et al, ³² 2002	UK	Male, 1557; female, 5702	48.7; 45.9	100; 0	125.5; 119.8	77.2; 73.7	46.3; 58.4 ^b	23.9; 23.4	<1 g ethanol/d, 23.0%/11.2%; 23.9%/20.3%	Mixed (vegan, lacto-ovo), current
Lu et al, ³³ 2000	Taiwan	Male, 53; female, 56	38.0; 39.2	100; 0	106.1; 99.5	70.0; 64.8	None	21.9; 21.2	Participants using >10 g/d excluded	Mixed (vegan, lacto), >2 y
Famodu et al, ³⁴ 1998	Nigeria	76	48.6	NR	110.1	76.1	NR	28.8	Veg group abstinent	Mixed (vegan, lacto-ovo, semi), current
Harman et al, ³⁵ 1998	New Zealand	Male, 23; female, 24	44.7; 41.0	100; 0	123.5; 115.0	77.4; 72.9	NR	25.2; 24.7	Excluded	Mixed (vegan, lacto), current
Williams, ³⁶ 1997	US	Male, 7253; female, 1989	45.9; 40.3	100; 0	121.7; 112.9	77.1; 71.7	NR	23.8; 21.2	39.1/85.6 mL; 26.3/52.8 mL	Mixed (vegan, lacto), current
Wyatt et al, ³⁷ 1995	Mexico	72	39.0	NR	115.0	78.5	None	24.4	1%/89%	Lacto-ovo, >1 y (mean, 5.1 y)
Melby et al, ³⁸ 1994	US	167	47.5	26.3	119.2	77.7	26.7	28.1	0/2	Mixed (vegetarian, semi), current

(continued)

Observational Studies

Thirty-two observational studies were identified (Table 2).¹⁶⁻⁴⁷ These studies included 21 604 participants (median sample size, 152; range, 20-9242) with a mean age of 46.6 years (range, 28.8-68.4 years). Each of the 32 observational studies used cross-sectional designs. As shown in Table 2, several participants in 5 observational studies^{17,28,32,38,40} used anti-hypertensive medication. Because pooled effects were not reported, male and female subgroups (10 studies)* and racial

*References 24, 25, 32, 33, 35, 36, 41, 44-46

subgroups (1 study)⁴⁰ were included in the subgroup analyses (Table 2). In 22 of these studies, participants had been following vegetarian diets for more than 1 year.† Five studies focused on vegan diets,^{23,27,31,39,41} 2 on lacto-vegetarian diets,^{18,24} 10 on lacto-ovo-vegetarian diets,‡ and 15 on mixed diet types (vegan, lacto, lacto-ovo, pescos, and/or semi-vegetarian).§ In 20 studies, diets were assessed by using

†References 16-20, 22-24, 26-31, 33, 37, 39-43, 45

‡References 16, 19, 20, 22, 26, 29, 37, 43, 44, 47

§References 17, 21, 25, 28, 30, 32-36, 38, 40, 42, 45, 46

Table 2. Designs and Population Characteristics of Observational Cross-Sectional Studies of Vegetarian Diets and BP (continued)

Source	Country	No. of Participants	Mean Age, y	Male Sex, %	Mean BP, mm Hg		Using Antihypertensive Medication, %	Mean BMI	Alcohol Intake (Veg/Cont)	Exposure, Duration of Exposure ^a
					Systolic	Diastolic				
Orlov et al, ³⁹ 1994	Finland	20	49.2	NR	131.3	78.9	NR	21.4	NR	Vegan, >1 y
Melby et al, ⁴⁰ 1989	US	Black, 114; white, 264	55.4; 52.5	21.9; 15.9	126.4; 114.8	74.5; 66.9	31.5; 12.7	28.9; 25.7	None; 1/2	Mixed (meatless), 20.6 y; 27.2 y
Sanders and Key, ⁴¹ 1987	UK	Male, 22; female, 22	31.5; 26.0	100; 0	115.0; 111.0	70.5; 68.5	None	21.5; 20.6	6/21 g/d; 0/1 g/d	Vegan, >1 y
Wiseman et al, ⁴² 1987	UK	52	34.4	48.1	113.5	73.1	None	NR	NR	Mixed (vegan, lacto), >3 y (mean, 13.4 y)
Ophir et al, ⁴³ 1983	Israel	196	60.9	51.0	136.4	82.3	NR	NR	Veg group abstinent	Lacto-ovo, >3 y (mean, 19 y)
Rouse et al, ⁴⁴ 1983	Australia	Male, 80; female, 100	32.8; 33.7	100; 0	117.9; 109.5	69.9; 67.0	None	23.6; 23.8	BP not related to past use of alcohol	Lacto-ovo, current
Burr et al, ⁴⁵ 1981	UK	Male, 111; female, 189	56.0; 53.6	100; 0	137.5; 140.4	84.6; 84.2	NR	24.0; 23.3	Weekly drinkers, 31%/50%; 23.2%/39.1%	Mixed (meat and fish less than once a month or never), >5 y
Haines et al, ⁴⁶ 1980	UK	Male, 236; female, 96	48.7; 49.5	100; 0	137.8; 137.7	87.1; 86.2	NR	NR	NR	Mixed (vegan, lacto-ovo), current
Armstrong et al, ⁴⁷ 1979	Australia	204	50.6	33.2	145.0	89.9	NR	23.5	NR (partly adjusted)	Lacto-ovo, current


Abbreviations: BP, blood pressure; BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); Cont, control; NR, not reported; UK, United Kingdom; US, United States; Veg, vegetarian.

omitting all animal products or vegetarian diets that include some animal products as indicated by the terms *lacto* (dairy products), *ovo* (eggs), or *pesco* (fish).

^a Semivegetarian diets, defined as rarely including meat; vegan diets, defined as

^b Included nutritional supplements.


Figure 2. Pooled Systolic and Diastolic Blood Pressure (BP) Responses to Vegetarian Diets in Clinical Trials



Effects on systolic BP (A) and on diastolic BP (B) are depicted as squares; error bars indicate 95% CIs. Meta-analysis yielded pooled estimates of -4.8 mm Hg (95% CI, -6.6 to -3.1) for systolic BP and -2.2 mm Hg (-3.5 to -1.0) for diastolic

BP, which are depicted as blue diamonds. Vegan diets were defined as omitting all animal products; vegetarian diets may include some animal products as indicated by the terms *lacto* (dairy products) and *ovo* (eggs).

Figure 3. Pooled Systolic Blood Pressure (BP) Among Vegetarians in Observational Studies



Effects on systolic BP are depicted as squares; error bars indicate 95% CIs. Meta-analysis yielded a pooled estimate of -6.9 mm Hg (95% CI, -9.1 to -4.7) for systolic BP, which is depicted as a blue diamond. Arrows indicate that the

95% CI exceeds the left line. Vegan diets were defined as omitting all animal products; vegetarian diets may include some animal products as indicated by the terms *lacto* (dairy products) and *ovo* (eggs).

questionnaires, typically food frequency questionnaires^{17,29,32,34,35,38,40} or 24-hour diet recalls. || Interviews or self-report were used in 7 studies,^{18,19,24,28,36,43,47} and weighing methods were used for 1 study²¹; the means of dietary assessment were not reported in 4 studies.^{23,26,30,46} Of the 32 observational studies, 12 conducted repeated measurements of BP.# The adjusted factors in each study are shown in the Supplement (eTable 2).

Pooled Effects of Vegetarian Diets on BP

In the clinical trials, consumption of vegetarian diets was associated with a mean reduction in systolic BP (-4.8 mm Hg; 95% CI, -6.6 to -3.1; $P < .001$; $I^2 = 0$; $P = .45$ for heterogeneity) and diastolic BP (-2.2 mm Hg; 95% CI, -3.5 to -1.0;

$P < .001$; $I^2 = 0$; $P = .43$ for heterogeneity) compared with the consumption of omnivorous diets (Figure 2).


In the observational studies, consumption of vegetarian diets was associated with lower mean systolic BP (-6.9 mm Hg; 95% CI, -9.1 to -4.7; $P < .001$; $I^2 = 91.4$; $P < .001$ for heterogeneity) and diastolic BP (-4.7 mm Hg; 95% CI, -6.3 to -3.1; $P < .001$; $I^2 = 92.6$; $P < .001$ for heterogeneity) compared with consumption of omnivorous diets (Figure 3 and Figure 4).

Meta-regression

In the meta-regression investigating the sources of heterogeneity in the observational trials, the potential sources were sex (proportion of men) (β coefficient, -0.03; $P < .001$), baseline systolic BP (-0.13; $P = .003$), baseline diastolic BP (-0.30; $P < .001$), sample size (0.001; $P < .001$), and BMI (-0.46; $P = .02$).

|| References 16, 20, 22, 25, 27, 31, 33, 37, 39, 41, 42, 44, 45
#References 16-18, 24, 27, 28, 31, 34, 35, 38, 40, 44

Figure 4. Pooled Diastolic Blood Pressure (BP) Among Vegetarians in Observational Studies



Effects on diastolic BP are depicted as squares; error bars indicate 95% CIs. Meta-analysis yielded a pooled estimate of -4.7 mm Hg (95% CI, -6.3 to -3.1) for diastolic BP, which is depicted as a blue diamond. Arrows indicate that the

95% CI exceeds the left line. Vegan diets were defined as omitting all animal products; vegetarian diets may include some animal products as indicated by the terms *lacto* (dairy products) and *ovo* (eggs).

These factors were not significant in the meta-regression of clinical trials (data not shown). These results suggest that the association between vegetarian diets and lower BP in adults is stronger among men and those with higher baseline BP and BMI. The association is also stronger in studies with smaller sample sizes.

Subgroup Analysis

Pooled changes in BP associated with consumption of vegetarian diets in planned strata are summarized in the Supplement (eTables 3 and 4). In the clinical trials, no heterogeneity was found in any subgroup and the estimated effect sizes were very similar.

For observational studies, subgrouping reduced heterogeneity in most cases, and vegetarian diets were associated with lower BP regardless of subgroup, although effect sizes were attenuated in some groups. Lower systolic BP values were


reported in the predominantly (50%-99%) male subgroups compared with the 100% female subgroups. Lower systolic and diastolic BP were found in both BMI subgroups (<25 and ≥25), sample size subgroups (<100 and ≥100), and baseline BP subgroups (normal, prehypertension, and stage 1 hypertension).

In the one-study-removed analysis, results were largely unchanged, with BP differences between the vegetarian and comparison groups ranging from -5.3 to -3.5 mm Hg for systolic BP and -2.9 to -1.8 mm Hg for diastolic BP in clinical trials (all results, *P* < .05) and from -7.2 to -6.3 mm Hg for systolic BP and -5.0 to -4.3 mm Hg for diastolic BP for observational studies (all results, *P* < .001).

Publication Bias

For clinical trials, visual examination of the funnel plot revealed that smaller trials that reported small reductions in systolic BP were possibly overrepresented (Figure 5A). In the

Figure 5. Funnel Plot of Comparison of Weight and Differences in Mean Systolic Blood Pressure (BP) Associated With Consumption of Vegetarian Diets



Funnel plot of study weights against change in systolic blood pressure (BP) in clinical trials (A) and observational studies (B). Blood pressure results in individual studies are depicted as circles scattered around the pooled BP estimate. A trim-and-fill method indicated that 3 clinical trials and no

observational studies might have been missing owing to publication bias. After adjustment for putative missing data, the overall differences for systolic BP increased to -5.2 mm Hg (95% CI, -6.9 to -3.5) in clinical trials.

absence of publication bias, study results would be symmetrically represented about the mean effect size; our findings suggest that a few studies were missing in the bottom left side. This visual impression was confirmed by the Egger test ($P = .04$). The results of use of the trim-and-fill method suggest that 3 trials might have been missing such that their addition would change the overall effect on systolic BP to -5.2 mm Hg (95% CI, -6.9 to -3.5).

For observational studies, visual examination of the funnel plot revealed that larger trials that reported generous reductions in systolic BP were possibly overrepresented. Our findings suggest that a few studies were missing in the middle right side (Figure 5B). This visual impression was confirmed by the Egger test ($P < .001$). The results of the trim-and-fill method suggest that no study was missing.

Discussion

This meta-analysis of 7 controlled trials and 32 observational studies indicates that consumption of vegetarian diets is associated with lower BP compared with consumption of omnivorous diets. The meta-analysis indicates an overall difference in systolic BP of -4.8 mm Hg in controlled trials and -6.9 mm Hg in observational studies. For diastolic BP, the differences were -2.2 mm Hg in controlled trials and -4.7 mm Hg in observational studies. These effect sizes are similar to those observed with commonly recommended lifestyle modifications, such as adoption of a low-sodium diet⁴⁸ or a weight reduction of 5 kg,⁴⁹ and are approximately half the magnitude of those observed with pharmaceutical therapy, such as administration of angiotensin-converting enzyme inhibitors to individuals with hypertension.⁵⁰ According to Whelton et al,⁵¹ a reduction in systolic BP of 5 mm Hg would

be expected to result in a 7%, 9%, and 14% overall reduction in mortality due to all causes, coronary heart disease, and stroke, respectively.

The findings of the present study are consistent with those of a previous review of observational studies.⁵ They also accord with those of the Dietary Approaches to Stop Hypertension study,^{52,53} which was based on the observation that consumption of vegetarian diets was associated with a reduced risk of hypertension and found that a diet rich in vegetables and fruits, along with other dietary changes, reduced systolic BP and diastolic BP.

Specific diet and lifestyle factors are known to influence BP. Obesity, excessive sodium intake, and excessive alcohol use are associated with increased BP and risk of hypertension; potassium intake and physical activity are associated with lower BP.^{54,55} In addition, intake of unsaturated fat, protein, magnesium, and dietary fiber may be associated with differences in BP.⁵ The details provided in the studies included in the present review were insufficient to justify subgroup analyses that might have investigated the influence of these factors on the observed BP differences. Nonetheless, the following factors merit consideration as possible explanations for the observed associations. First, compared with omnivores, vegetarians typically have lower BMIs and a lower risk of obesity, which is mainly attributable to the lower energy density of the diet that results from higher fiber content and lower fat content.⁵⁶ Weight differences do not fully explain the observed BP differences, however, because studies controlling for body weight have demonstrated a BP-lowering effect of vegetarian diets.⁶ Second, potassium is abundant in vegetarian diets.⁵⁷ Meta-analyses^{58,59} of randomized clinical trials have reported that potassium supplementation decreases BP. It is hypothesized that a high potassium intake increases vasodilation and glomerular filtration rate while decreasing renin level,

renal sodium reabsorption, reactive oxygen species production, and platelet aggregation.⁶⁰ Third, some reports⁶¹ have suggested that vegetarian diets may be lower in sodium; however, others⁵⁷ have shown no clear differences in sodium intake between nonvegetarians and vegetarians. Fourth, some studies^{32,36,37,41} have reported that alcohol consumption is lower in vegetarian populations compared with the general population. However, of the 7 clinical trials included in our study, 5 were limited to participants with no more than modest alcohol consumption; their results are unlikely to be substantially affected by alcohol intake. Vegetarian diets are often proportionately lower in saturated fatty acids and richer in polyunsaturated fatty acids compared with omnivorous diets; both of these dietary characteristics are associated with lower BP.^{5,62,63} Consumption of vegetarian diets has also been associated with reduced blood viscosity, which may affect BP.⁶⁴ The consumption of vegetable protein has been shown to be inversely associated to BP.⁶⁵

The present meta-analysis has several strengths. First, the available clinical trials and observational studies provided a reasonably large overall sample size that fosters confidence in the findings as well as permitting subgroup analyses in specific population groups. Second, its focus on dietary patterns rather than on the use of dietary supplements or artificial dietary manipulations makes the findings easily applicable to general or clinical populations.

This review also has several limitations. First, although no heterogeneity existed among the controlled trials, heterogeneity was high among the observational studies.

Meta-regression and subgroup analyses showed that sex, baseline BP, sample size, and BMI may be key reasons for this heterogeneity. Nonetheless, lower BP was evident in all subgroups, although the differences were not significant for some subgroups. Second, this meta-analysis carried forward design limitations of the included studies. Most notable in this regard are small sample sizes and the fact that all observational studies used cross-sectional rather than prospective designs; however, the latter limitation is partially compensated for by the inclusion of several randomized clinical trials. Third, some of the observational studies did not adjust for lifestyle factors, such as alcohol intake or physical activity level. Finally, foods that make up vegetarian diets and the nutrient composition of the diets differ from person to person and from country to country. Further studies are needed to explore the relationships between specific foods and nutrients and BP. Nevertheless, the results of the meta-analysis of the controlled trials suggest a robust relationship between consumption of vegetarian diets and lower BP.

Conclusions

Consumption of vegetarian diets is associated with lower BP. Further studies are required to clarify which types of vegetarian diets are most strongly associated with lower BP. Research into the implementation of such diets, either as public health initiatives aiming at prevention of hypertension or in clinical settings, would also be of great potential value.

ARTICLE INFORMATION

Accepted for Publication: December 8, 2013.

Published Online: February 24, 2014.
doi:10.1001/jamainternmed.2013.14547.

Author Affiliations: Department of Preventive Medicine and Epidemiologic Informatics, National Cerebral and Cardiovascular Center, Osaka, Japan (Yokoyama, Takegami, Miyamoto); Japan Society for the Promotion of Science, Tokyo, Japan (Yokoyama); Physicians Committee for Responsible Medicine, Washington, DC (Yokoyama, Barnard); Division of Evidence-Based Medicine and Risk Analysis, Preventive Medicine and Epidemiology, National Cerebral and Cardiovascular Center, Osaka, Japan (Nishimura); Department of Nephrology, School of Medicine, Fujita Health University, Aichi, Japan (Nishimura); Department of Medicine, George Washington University School of Medicine and Health Sciences, Washington, DC (Barnard); Department of Healthcare Epidemiology, Kyoto University School of Medicine and Public Health, Kyoto, Japan (Takegami); Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, Osaka, Japan (Watanabe, Miyamoto); Department of Epidemiology, University of Pittsburgh, Pittsburgh, Pennsylvania (Sekikawa); Department of Preventive Medicine and Public Health, School of Medicine, Keio University, Tokyo, Japan (Okamura).

Author Contributions: Dr Yokoyama had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Yokoyama, Nishimura, Barnard, Okamura, Miyamoto.

Acquisition of data: Yokoyama.

Analysis and interpretation of data: Yokoyama, Nishimura, Barnard, Takegami, Watanabe, Sekikawa.

Drafting of the manuscript: Yokoyama, Nishimura, Barnard, Miyamoto.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Yokoyama, Nishimura.

Obtained funding: Yokoyama.

Administrative, technical, or material support: Yokoyama, Takegami, Miyamoto.

Study supervision: All authors.

Conflict of Interest Disclosures: None reported.

Funding/Support: Financial support for this study was provided by a grant-in-aid for Japan Society for the Promotion of Science Fellows grant 23-10883.

Role of the Sponsor: The funding source had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: Richard Holubkov, PhD, Division of Critical Care, Department of Pediatrics, University of Utah School of Medicine, provided statistical advice, and T. Colin Campbell, PhD, Nutritional Biochemistry, Division of Nutritional Sciences, Cornell University, provided conceptual advice. Dr Holubkov received compensation for his assistance.

REFERENCES

1. Chobanian AV, Bakris GL, Black HR, et al; National Heart, Lung, and Blood Institute Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure; National High Blood Pressure Education Program Coordinating Committee. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*. 2003;289(19):2560-2572.
2. Lewington S, Clarke R, Qizilbash N, Peto R, Collins R; Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 2002;360(9349):1903-1913.
3. Chobanian AV, Bakris GL, Black HR, et al; Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure; National Heart, Lung, and Blood Institute; National High Blood Pressure Education Program Coordinating Committee. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension*. 2003;42(6):1206-1252.
4. Hu FB. Plant-based foods and prevention of cardiovascular disease: an overview. *Am J Clin Nutr*. 2003;78(3)(suppl):544S-551S.
5. Berkow SE, Barnard ND. Blood pressure regulation and vegetarian diets. *Nutr Rev*. 2005;63(1):1-8.

6. Rouse IL, Beilin LJ, Armstrong BK, Vandongen R. Blood-pressure-lowering effect of a vegetarian diet: controlled trial in normotensive subjects. *Lancet*. 1983;321(8314-8315):5-10.
7. Margetts BM, Beilin LJ, Vandongen R, Armstrong BK. Vegetarian diet in mild hypertension: a randomised controlled trial. *Br Med J (Clin Res Ed)*. 1986;293(6560):1468-1471.
8. Kestin M, Rouse IL, Correll RA, Nestel PJ. Cardiovascular disease risk factors in free-living men: comparison of two prudent diets, one based on lactoovovegetarianism and the other allowing lean meat. *Am J Clin Nutr*. 1989;50(2):280-287.
9. de Mello VD, Zelmanovitz T, Perassolo MS, Azevedo MJ, Gross JL. Withdrawal of red meat from the usual diet reduces albuminuria and improves serum fatty acid profile in type 2 diabetes patients with macroalbuminuria. *Am J Clin Nutr*. 2006;83(5):1032-1038.
10. Follmann D, Elliott P, Suh LI, Cutler J. Variance imputation for overviews of clinical trials with continuous response. *J Clin Epidemiol*. 1992;45:769-773.
11. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327(7414):557-560.
12. Ferdowsian HR, Barnard ND, Hoover VJ, et al. A multicomponent intervention reduces body weight and cardiovascular risk at a GEICO corporate site. *Am J Health Promot*. 2010;24(6):384-387.
13. Nicholson AS, Sklar M, Barnard ND, Gore S, Sullivan R, Browning S. Toward improved management of NIDDM: a randomized, controlled, pilot intervention using a lowfat, vegetarian diet. *Prev Med*. 1999;29(2):87-91.
14. Sciarra SE, Strahan MT, Beilin LJ, Burke V, Rogers P, Rouse IL. Biochemical and neurohormonal responses to the introduction of a lacto-ovovegetarian diet. *J Hypertens*. 1993;11(8):849-860.
15. Hakala P, Karvetti RL. Weight reduction on lacto-vegetarian and mixed diets: changes in weight, nutrient intake, skinfold thicknesses and blood pressure. *Eur J Clin Nutr*. 1989;43(6):421-430.
16. Kim MH, Bae YJ. Postmenopausal vegetarians' low serum ferritin level may reduce the risk for metabolic syndrome. *Biol Trace Elem Res*. 2012;149(1):34-41.
17. Pettersen BJ, Anousheh R, Fan J, Jaceldo-Siegl K, Fraser GE. Vegetarian diets and blood pressure among white subjects: results from the Adventist Health Study-2 (AHS-2). *Public Health Nutr*. 2012;15(10):1909-1916.
18. Yang SY, Li XJ, Zhang W, et al. Chinese lacto-vegetarian diet exerts favorable effects on metabolic parameters, intima-media thickness, and cardiovascular risks in healthy men. *Nutr Clin Pract*. 2012;27(3):392-398.
19. Chen CW, Lin CT, Lin YL, Lin TK, Lin CL. Taiwanese female vegetarians have lower lipoprotein-associated phospholipase A2 compared with omnivores. *Yonsei Med J*. 2011;52(1):13-19.
20. Fernandes Dourado K, de Arruda Câmara e Siqueira Campos F, Sakugava Shinohara NK. Relation between dietary and circulating lipids in lacto-ovo vegetarians. *Nutr Hosp*. 2011;26(5):959-964.
21. Rodenas S, Sánchez-Muniz FJ, Bastida S, Sevillano MI, Larrea Marín T, González-Muñoz MJ. Blood pressure of omnivorous and semi-vegetarian postmenopausal women and their relationship with dietary and hair concentrations of essential and toxic metals. *Nutr Hosp*. 2011;26(4):874-883.
22. Yang SY, Zhang HJ, Sun SY, et al. Relationship of carotid intima-media thickness and duration of vegetarian diet in Chinese male vegetarians. *Nutr Metab (Lond)*. 2011;8(1):63. doi:10.1186/1743-7075-8-63.
23. Lin CK, Lin DJ, Yen CH, et al. Comparison of renal function and other health outcomes in vegetarians versus omnivores in Taiwan. *J Health Popul Nutr*. 2010;28(5):470-475.
24. Pitla S, Nagala B. Gender-related differences in the relationship between plasma homocysteine, anthropometric and conventional biochemical coronary heart disease risk factors in middle-aged Indians. *Ann Nutr Metab*. 2009;54(1):1-6.
25. Nakamoto K, Watanabe S, Kudo H, Tanaka A. Nutritional characteristics of middle-aged Japanese vegetarians. *J Atheroscler Thromb*. 2008;15(3):122-129.
26. Slavicek J, Kittnar O, Fraser GE, et al. Lifestyle decreases risk factors for cardiovascular diseases. *Cent Eur J Public Health*. 2008;16(4):161-164.
27. Fontana L, Meyer TE, Klein S, Holloszy JO. Long-term low-calorie low-protein vegan diet and endurance exercise are associated with low cardiometabolic risk. *Rejuvenation Res*. 2007;10(2):225-234.
28. Teixeira R de C, Molina M del C, Zandonade E, Mill JG. Cardiovascular risk in vegetarians and omnivores: a comparative study. *Arq Bras Cardiol*. 2007;89(4):237-244.
29. Sebeková K, Boor P, Valachovicová M, et al. Association of metabolic syndrome risk factors with selected markers of oxidative status and microinflammation in healthy omnivores and vegetarians. *Mol Nutr Food Res*. 2006;50(9):858-868.
30. Su TC, Jeng JS, Wang JD, et al. Homocysteine, circulating vascular cell adhesion molecule and carotid atherosclerosis in postmenopausal vegetarian women and omnivores. *Atherosclerosis*. 2006;184(2):356-362.
31. Goff LM, Bell JD, So PW, Dornhorst A, Frost GS. Veganism and its relationship with insulin resistance and intramyocellular lipid. *Eur J Clin Nutr*. 2005;59(2):291-298.
32. Appleby PN, Davey GK, Key TJ. Hypertension and blood pressure among meat eaters, fish eaters, vegetarians and vegans in EPIC-Oxford. *Public Health Nutr*. 2002;5(5):645-654.
33. Lu SC, Wu WH, Lee CA, Chou HF, Lee HR, Huang PC. LDL of Taiwanese vegetarians are less oxidizable than those of omnivores. *J Nutr*. 2000;130(6):1591-1596.
34. Famodu AA, Osilei O, Makinde YO, Osonuga OA. Blood pressure and blood lipid levels among vegetarian, semi-vegetarian, and non-vegetarian native Africans. *Clin Biochem*. 1998;31(7):545-549.
35. Harman SK, Parnell WR. The nutritional health of New Zealand vegetarian and non-vegetarian Seventh-day Adventists: selected vitamin, mineral and lipid levels. *N Z Med J*. 1998;111(1062):91-94.
36. Williams PT. Interactive effects of exercise, alcohol, and vegetarian diet on coronary artery disease risk factors in 9242 runners: the National Runners' Health Study. *Am J Clin Nutr*. 1997;66(5):1197-1206.
37. Wyatt CJ, Velazquez C, Grijalva I, Valencia ME. Dietary-intake of sodium, potassium and blood-pressure in lacto-ovo-vegetarians. *Nutr Res*. 1995;15(6):819-830. doi:10.1016/0271-5317(95)00048-N.
38. Melby CL, Toohey ML, Cebrick J. Blood pressure and blood lipids among vegetarian, semivegetarian, and nonvegetarian African Americans. *Am J Clin Nutr*. 1994;59(1):103-109.
39. Orlov SN, Agren JJ, Hänninen OO, et al. Univalent cation fluxes in human erythrocytes from individuals with low or normal sodium intake. *J Cardiovasc Risk*. 1994;1(3):249-254.
40. Melby CL, Goldflies DG, Hyner GC, Lyle RM. Relation between vegetarian/nonvegetarian diets and blood pressure in black and white adults. *Am J Public Health*. 1989;79(9):1283-1288.
41. Sanders TA, Key TJ. Blood pressure, plasma renin activity and aldosterone concentrations in vegans and omnivore controls. *Hum Nutr Appl Nutr*. 1987;41(3):204-211.
42. Wiseman MJ, Hunt R, Goodwin A, Gross JL, Keen H, Viberti GC. Dietary composition and renal function in healthy subjects. *Nephron*. 1987;46(1):37-42.
43. Ophir O, Peer G, Gilad J, Blum M, Aviram A. Low blood pressure in vegetarians: the possible role of potassium. *Am J Clin Nutr*. 1983;37(5):755-762.
44. Rouse IL, Armstrong BK, Beilin LJ. The relationship of blood pressure to diet and lifestyle in two religious populations. *J Hypertens*. 1983;1(1):65-71.
45. Burr ML, Bates CJ, Fehily AM, St Leger AS. Plasma cholesterol and blood pressure in vegetarians. *J Hum Nutr*. 1981;35(6):437-441.
46. Haines AP, Chakrabarti R, Fisher D, Meade TW, North WR, Stirling Y. Haemostatic variables in vegetarians and non-vegetarians. *Thromb Res*. 1980;19(1-2):139-148.
47. Armstrong B, Clarke H, Martin C, Ward W, Norman N, Masarei J. Urinary sodium and blood pressure in vegetarians. *Am J Clin Nutr*. 1979;32(12):2472-2476.
48. Graudal NA, Hubeck-Graudal T, Jürgens G. Effects of low-sodium diet vs. high-sodium diet on blood pressure, renin, aldosterone, catecholamines, cholesterol, and triglyceride (Cochrane Review). *Am J Hypertens*. 2012;25(1):1-15.
49. Neter JE, Stam BE, Kok FJ, Grobbee DE, Geleijnse JM. Influence of weight reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension*. 2003;42(5):878-884.
50. Brugts JJ, Ninomiya T, Boersma E, et al. The consistency of the treatment effect of an ACE-inhibitor based treatment regimen in patients with vascular disease or high risk of vascular disease: a combined analysis of individual data of ADVANCE, EUROPA, and PROGRESS trials. *Eur Heart J*. 2009;30(11):1385-1394.
51. Whelton PK, He J, Appel LJ, et al; National High Blood Pressure Education Program Coordinating Committee. Primary prevention of hypertension: clinical and public health advisory from the National High Blood Pressure Education Program. *JAMA*. 2002;288(15):1882-1888.

52. Appel LJ, Moore TJ, Obarzanek E, et al; DASH Collaborative Research Group. A clinical trial of the effects of dietary patterns on blood pressure. *N Engl J Med*. 1997;336(16):1117-1124.
53. Sacks FM, Obarzanek E, Windhauser MM, et al. Rationale and design of the Dietary Approaches to Stop Hypertension trial (DASH): a multicenter controlled-feeding study of dietary patterns to lower blood pressure. *Ann Epidemiol*. 1995;5(2):108-118.
54. Koliaki C, Katsilambros N. Dietary sodium, potassium, and alcohol: key players in the pathophysiology, prevention, and treatment of human hypertension. *Nutr Rev*. 2013;71(6):402-411.
55. Frisoli TM, Schmieder RE, Grodzicki T, Messerli FH. Beyond salt: lifestyle modifications and blood pressure. *Eur Heart J*. 2011;32(24):3081-3087.
56. Berkow SE, Barnard N. Vegetarian diets and weight status. *Nutr Rev*. 2006;64(4):175-188.
57. Rizzo NS, Jaceldo-Siegl K, Sabate J, Fraser GE. Nutrient profiles of vegetarian and nonvegetarian dietary patterns. *J Acad Nutr Diet*. 2013;113(12):1610-1619.
58. Whelton PK, He J, Cutler JA, et al. Effects of oral potassium on blood pressure: meta-analysis of randomized controlled clinical trials. *JAMA*. 1997;277(20):1624-1632.
59. Aburto NJ, Hanson S, Gutierrez H, Hooper L, Elliott P, Cappuccio FP. Effect of increased potassium intake on cardiovascular risk factors and disease: systematic review and meta-analyses. *BMJ*. 2013;346:f1378. doi:10.1136/bmj.f1378.
60. McDonough AA, Nguyen MT. How does potassium supplementation lower blood pressure? *Am J Physiol Renal Physiol*. 2012;302(9):F1224-F1225.
61. Larsson CL, Johansson GK. Dietary intake and nutritional status of young vegans and omnivores in Sweden. *Am J Clin Nutr*. 2002;76(1):100-106.
62. Iacono JM, Dougherty RM. Effects of polyunsaturated fats on blood pressure. *Annu Rev Nutr*. 1993;13:243-260.
63. Stamler J, Caggiula A, Grandits GA, Kjelsberg M, Cutler JA. Relationship to blood pressure of combinations of dietary macronutrients: findings of the Multiple Risk Factor Intervention Trial (MRFIT). *Circulation*. 1996;94(10):2417-2423.
64. Ernst E, Pietsch L, Matrai A, Eisenberg J. Blood rheology in vegetarians. *Br J Nutr*. 1986;56(3):555-560.
65. Elliott P, Stamler J, Dyer AR, et al. Association between protein intake and blood pressure: the INTERMAP Study. *Arch Intern Med*. 2006;166(1):79-87.