

Plant-based diets for children as a means of improving adult cardiometabolic health

Małgorzata A. Desmond, Jakub Sobiecki, Mary Fewtrell, and Jonathan C.K. Wells

Cardiovascular disease (CVD) is the largest contributor to global mortality, and this trend is expected to continue. Although mortality rates have been falling, adverse developments in obesity and diabetes threaten to reverse this. It has been estimated that the only viable strategy to reduce the epidemic is to focus on population-wide risk factor reduction. Primordial prevention, a strategy aimed at avoiding the development of risk factors before the disease onset, has been shown to reduce the CVD epidemic substantially. Plant-based diets appear beneficial for prevention of cardiometabolic diseases, with adult vegetarians and vegans having lower CVD risk than omnivores. Atherosclerosis starts in childhood and progresses in relation to classical CVD risk factors, which, along with dietary habits, track to adulthood. Based on this evidence, it is proposed that plant-based diets in childhood could promote cardiometabolic health in adults and thereby reduce CVD and promote longevity and health. However, the need for additional research to establish the safety of predominantly or exclusively plant-based diets in children is noted.

INTRODUCTION

Cardiovascular disease (CVD) remains the number one cause of premature mortality in the world.¹ In high-income countries, it also contributes the highest percentage of ill health in adults.² In most industrialized countries, CVD death rates have been declining since the late 1970s.³ However, 2 factors threaten to reverse this trend today: aging of the population (ie, increased absolute numbers of those aged > 70 and > 80 y), which increases the lifetime exposure to the risk factors; and global increases in the prevalence of obesity and diabetes⁴ and, in some countries, in mean blood pressure and smoking prevalence.⁵ These trends may be beginning to cancel out the health gains linked to declines in other risk factors and better care and treatment achieved in recent decades as recently shown for younger adults in

the United Kingdom, United States, and Australia.^{4–6} The consequence of these antagonistic trends might be an increase in the actual burden of CVD.⁴

It has been postulated that the only strategy capable of substantially reducing the CVD burden in the current scenario is to focus on population-wide reduction of major CVD risk factors, particularly targeting cholesterol levels and blood pressure. This approach has been shown to be more effective than focusing on pharmacological intervention in high-risk individuals.⁷ The potential impact of implementing risk-factor reduction policies focused on diet and lifestyle in 9 European countries has been recently quantified, and it was estimated that it would result in up to 29.1% fewer CVD deaths by 2020.⁸

However, there is increasing evidence that cardiometabolic risk in adulthood is partially determined by

Affiliation: M.A. Desmond, M. Fewtrell, and J.C.K. Wells are with the UCL Great Ormond Street Institute of Child Health, University College London, London, United Kingdom; M.A. Desmond and J. Sobiecki are with the Department of Paediatrics, Nutrition and Metabolic Diseases, Children's Memorial Health Institute, Warsaw, Poland. J. Sobiecki is with the School of Public Health, Imperial College, London, United Kingdom.

Correspondence: M. Desmond, Childhood Nutrition Research Centre, Population, Policy and Practice, UCL Great Ormond St Institute of Child Health, 30 Guilford St, London, WC1N 1EH, UK. E-mail: malgorzata.desmond.11@ucl.ac.uk

Key words: children, cardiovascular disease, cardiovascular disease risk factors, primordial prevention, vegetarian diets.

© The Author(s) 2018. Published by Oxford University Press on behalf of the International Life Sciences Institute. All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.

the same risk profile in childhood.⁹ Therefore, interventions that target risk factors in young people could potentially be key aspects of preventive strategies in the long-term.

Diet is well established to be one of the most important factors affecting cardiovascular risk in adults.¹⁰ A number of axes of dietary variability have been linked with cardiometabolic risk, including the consumption of saturated and trans-fats, sodium, fruits and vegetables, whole grains, fish, and nuts.¹⁰ However, there is growing interest in one particular axis that may also relate to many of the others: namely, the proportion of the diet obtained from plants. Plant-based diets include vegan diets that exclude products of animal origin, vegetarian diets that rule out meat and fish consumption, and also dietary patterns predominantly based on plant foods with undefined and individually determined levels of animal product consumption. For the purpose of this review, vegetarian and vegan diets will be jointly defined as meat-free diets.

Beyond their potential health benefits, plant-based diets are also considered critical for long-term planetary sustainability,¹¹ and they are furthermore chosen for ethical reasons by those who sympathize with animal welfare movements.¹² It appears that interest in meat-free diets is growing in many countries¹³; however exact estimates of these trends are yet to be produced. These issues collectively justify greater research into how plant-based diets might be adapted to simultaneously promote health and decrease ecological damage.

In adults, dietary patterns emphasizing the intake of unprocessed foods of plant origin seem to confer substantial protection against CVD. In particular, consumption of fruit and vegetables, whole grains, nuts, and legumes has been linked inversely with CVD risk in a dose-dependent fashion.¹⁰ In this context, it does not seem surprising that adults systematically pursuing vegetarian and vegan diets have a more favorable profile of CVD risk factors,^{14,15} translating to lower prevalence of ischemic heart disease (IHD; the predominant form of CVD in Western populations) risk,^{16,17} than otherwise similar nonvegetarians, with this difference primarily attributed to diet.¹⁸ However, a note of caution is that studies are inconsistent regarding whether the rate of mortality from IHD differs between vegetarians/vegans and omnivores, with one study showing 26%–34% reductions in risk for different vegetarian diets¹⁷ but another study found no difference.¹⁹ Moreover, there is currently no evidence that vegetarians/vegans have lower rates of cerebrovascular disease.^{17,19,20}

Some of this inconsistency in the association of diet and mortality risk could potentially be explained by differences in the uptake of treatment and other health services between vegetarians and nonvegetarians or

differences in the duration of consuming the vegetarian diet. Another possibility is that the cardioprotective effects of plant-based diets relating to decreased IHD risk may be counterbalanced by a lack of beneficial effects on non-IHD CVD risk. Further research is required to improve understanding of these scenarios.

Although the overt manifestation of CVD occurs primarily in adult life, the origins of atherosclerosis start well before,²¹ and the atherogenic process is influenced by measurable risk factors. So far, research on childhood cardiometabolic risk has placed greatest emphasis on markers of growth and nutritional status, including birth weight,^{22,23} childhood body mass index (BMI),^{24,25} and catch-up growth.²⁶ However, these childhood risk factors are potentially influenced by dietary composition, and healthy childhood dietary patterns have been found to be associated with lower adulthood CVD risk.²⁷ This might be mediated via beneficial effects of prudent diets on the CVD risk factor profile in childhood other than just BMI.

This article reviews substantial evidence on 1) the association between meat-free diets and CVD risk in adults, 2) the developmental origins of atherosclerosis, and 3) the tendency for childhood cardiometabolic risk factors to track into adulthood. This generates a new hypothesis that meat-free or plant-based diets in childhood, through their effects on blood biochemistry and other cardiovascular risk factors, offer a novel opportunity to promote a healthy childhood trajectory toward adult cardiometabolic health.

PLANT-BASED DIETS AND ADULT CARDIOVASCULAR DISEASE RISK

Diet represents a key modifiable direct risk factor for CVD, and it also impacts other components of risk, including obesity, unhealthy lipid profile, hypertension, raised blood glucose, and even physical activity level. Foods of plant origin have the most established protective effect on CVD.¹⁰

As previously noted, vegetarian diets may decrease the risk of IHD, which is likely to be at least partly mediated via classical CVD risk factors. Descriptive analysis of cardiovascular profile of vegetarians may thus help elucidate mechanisms through which plant-based diets exert their cardioprotective effects.

First, vegetarians and vegans have lower levels of body fat, mainly characterized by BMI. A recent systematic review that incorporated meta-analysis of 71 cross-sectional studies that examined the effect of a vegetarian diet and 19 cross-sectional studies that examined the effect of a vegan diet showed that these diets are associated with a BMI that is 1.49 kg/m² (95% confidence interval [CI], −1.72 to −1.25) and

1.72 kg/m² (95%CI, −2.21 to −1.22) lower than that of the reference omnivorous population, respectively.²⁰ These lower BMI values are attributed to the lower energy density of diets high in fruits and vegetables and the satiating effect of increased fiber intakes.²⁸ The association of obesity with CVD risk is mediated by various other risk factors,²⁹ which also show a better profile among people following vegetarian diets.

Second, vegetarians, especially vegans, have lower levels of total and non-high-density lipoprotein cholesterol (HDL-C). A recent systematic review with meta-analysis of observational studies showed estimates of effect size ranging from reductions of 0.72 mmol/L (95%CI, −0.8 to −0.64) in total cholesterol (TC) and 0.55 mmol/L (95%CI, −0.62 to −0.47) in low-density lipoprotein cholesterol (LDL-C) associated with vegetarian diets compared with omnivore diets (based on 64 and 46 cross-sectional studies, respectively) to reductions of 0.80 mmol/L (95%CI, −0.90 to −0.70) in TC and 0.59 mmol/L (95%CI, −0.77 to −0.40) in LDL-C associated with vegan diets compared with omnivore diets (based on 19 and 13 cross-sectional studies, respectively).²⁰ Both total and non-HDL cholesterol concentrations tend to be highest in meat eaters and lowest in vegans, with vegetarians having intermediate values.³⁰ In another systematic review and meta-analysis of 11 randomized clinical trials (7 included a vegan diet, 2 included a lacto-ovo vegetarian diet, 2 included a lacto vegetarian diet), Wang et al.¹⁴ showed that vegetarian diets were associated with lower TC, LDL-C, and non-HDL-C, with pooled estimated effects of −0.36 mmol/L (95%CI, −0.55 to −0.17), −0.34 mmol/L (95%CI, −0.57 to −0.11), and −0.30 mmol/L (95%CI, −0.50 to −0.10), respectively. A 1-mmol/L reduction in TC and LDL-C levels results in a 26.6%–29.5% decrease in risk for any CVD-related event.³¹ Therefore, the average reductions of TC and LDL-C concentrations following a vegetarian diet would correspond to a decrease in CVD risk of approximately 9.0%–10.6%.¹⁴

Mechanistically, vegetarian diets may reduce blood cholesterol concentrations due to their lower content of saturated fat, total fat, and cholesterol³² and higher intakes of dietary fiber and numerous phytochemicals, all of which have been linked to lower blood lipids.³³

Third, adults on plant-based diets have lower systolic blood pressure (SBP) and diastolic blood pressure (DBP), and lower risk of hypertension compared with meat eaters. In the cross-sectional analysis of a subset of 592 black women and men enrolled in the Adventist Health Study 2 (25% vegetarian and vegan; 75% nonvegetarian) the risk of hypertension varied among dietary groups and was lowest for vegans and highest for omnivores. The relative risk (RR) in comparison with omnivores was 0.37 (95%CI, 0.19–0.74) and 0.57 (95%CI,

0.36–0.92) for vegans and vegetarians, respectively, in a model adjusted for age, sex, and physical activity.³⁴ In a matched cohort study of 4109 Taiwanese nonsmokers, where each vegetarian was matched with 5 nonvegetarians by age, sex, and study site, vegetarians had a 28% lower risk (RR = 0.72; 95%CI, 0.55–0.86) for hypertension adjusting for age, sex, C-reactive protein, waist circumference, and fasting glucose.³⁵

Similarly, a systematic review and meta-analysis of controlled clinical trials and observational studies showed a reduction in mean SBP (−4.8 mm Hg; 95%CI, −6.6 to −3.1) and DBP (−2.2 mm Hg; 95%CI, −3.5 to −1.0) after application of a vegetarian diet compared with the consumption of omnivorous diets (7 controlled trials with a total of 311 participants; mean age, 44.5 y) and lower mean SBP (−6.9 mm; 95%CI, −9.1 to −4.7) and DBP (−4.7 mm Hg; 95%CI, −6.3 to −3.1) associated with consumption of a vegetarian diet compared with an omnivorous diet (32 observational studies with a total of 21 604 participants; mean age, 46.6 y).¹⁵

Mechanistically, several dietary factors in plant-based diets, other than those affecting BMI, may account for the effects on BP. Vegetarians have higher fiber intakes,³⁶ which have been shown to lower blood pressure.³⁷ Other factors, including higher plant protein³⁸ and potassium intakes^{39,40} and lower heme iron intake,⁴¹ may improve blood pressure regulation. Potential mechanisms include baroreceptor sensitivity, direct vasodilatory effects, and changes in catecholamine and renin-angiotensin-aldosterone metabolism,⁴² along with changes in blood viscosity.⁴³

Every 10 mm Hg reduction in SBP reduces the risk of major CVD events (RR = 0.80; 95%CI, 0.77–0.83), coronary heart disease (RR = 0.83; 95%CI, 0.78–0.88), and stroke (RR = 0.73; 95%CI, 0.68–0.77).⁴⁴

Fourth, plant-based diets are associated with lower blood glucose levels, insulin resistance, and diabetes risk.^{45,46} In a systematic review and meta-analysis of observational studies, which included 27 studies with a total of 2256 vegetarian and 2192 nonvegetarian participants and 4 studies with a total of 83 vegans and 125 omnivores, plant-based diets were associated with lower blood glucose levels (vegetarians: −5.08 mg/dL, 95%CI, −5.98 to −4.19; vegans: 6.39 mg/dL, 95%CI, −12.35 to −0.41).²⁰ Observational studies further show that the prevalence of type 2 diabetes is 1.6–2 times lower in vegetarians than in omnivores, even after controlling for body weight.⁴⁷

Likewise, clinical interventions in individuals with type 2 diabetes have demonstrated that adopting a vegetarian diet leads to a greater reduction in fasting plasma glucose, HbA1c, and hypoglycemic medication compared with a conventional hypocaloric diet.^{48,49} These results confirm substantial protective effects of

plant-based diets,^{50–53} along with the avoidance of meat,⁵⁴ on glycemic control and risk of diabetes. Mechanistically, plant-based diets may confer protective effects on diabetes risk through caloric restriction; reduced intake of saturated fatty acids; high intake of polyunsaturated and monounsaturated fatty acids; low glycemic index; increased intake of fiber; higher intake of nonheme iron and reduction in iron stores; increased intake of antioxidants, vitamins, and micronutrients; high intake of vegetable instead of animal protein; and high intake of plant sterols and prebiotics. All of these have been shown to have a positive effect on diabetes prevention.⁴⁷

Diabetes confers an approximately 2-fold excess risk for CVD, independent of other conventional risk factors. In people without diabetes, fasting blood glucose concentration is modestly and nonlinearly associated with risk of vascular disease.⁵⁵

Finally, lower levels of C-reactive protein (CRP) have been reported in adult vegetarians (0.77 mg/L, standard error [SE], 1.29 for vegetarians; 1.30 mg/L, SE, 1.38 for matched omnivores; $P < 0.01$),⁵⁶ and a decrease in CRP was observed in adults adopting vegan diets (−28.2%; SE, 10.8%; $P = 0.02$).⁵⁷ Some studies have shown that healthy adult lactovegetarians have lower carotid intima media thickness (IMT),⁵⁸ with the advantage related to the duration of consumption of the vegetarian diet,⁵⁹ but 1 study showed no such difference.⁶⁰

Altogether, a lower prevalence of cardiometabolic risk factors among adults following plant-based diets is likely to be the primary reason why they have an approximately 25% lower risk of developing IHD.^{16,20} The overall pattern in which vegetarian or vegan diets affect IHD risk factors and incidence is illustrated in Figure 1. The effects of meat-free diets on cardiovascular risk factors are summarized in Table 1^{14,15,20,34,35,45,46,48,56,57}. However, although the main burden of CVD morbidity and mortality occurs in adults, there is compelling evidence that CVD risk is strongly shaped by experience at earlier ages.

ATHEROSCLEROSIS STARTS IN CHILDHOOD

Atherosclerosis comprises arterial lesions that are a fundamental component of adult CVD. These lesions develop over time, starting with the relatively harmless accumulation of lipid-filled macrophages, called fatty streaks, and progressing to more advanced stages where the streaks are raised and vulnerable to rupture, manifesting as fibrous and calcified plaques.⁶¹ Histological studies indicate that the clinically consequential lesions develop from these initially harmless changes in symptomatic individuals.⁶²




Figure 1 The overall pattern by which vegetarian/vegan diets affect ischemic heart disease risk factors and incidence. Abbreviations: CRP, C-reactive protein; IHD, ischemic heart disease; MUFA, monounsaturated fatty acid; PUFA, polyunsaturated fatty acid.

Early autopsy studies first suggested that atherosclerosis begins early in life. Enos et al.⁶³ showed gross evidence of coronary atherosclerosis among autopsies of 77.3% of US soldiers (average age, 22 years) killed in the Korean war.³

Around the same time, Holman et al.²¹ demonstrated the presence of fatty streaks in the aortas of children as young as 3 years. Subsequently, McNamara et al.⁶⁴ reported atherosclerosis in 45% and severe coronary atherosclerosis in 5% of 105 autopsies of US soldiers killed in Vietnam.

More recently, the Pathobiological Determinants of Atherosclerosis in Youth (PDAY) study described the emergence of clinically consequential atherosclerotic lesions in a large sample of autopsied persons aged 15–34 years who died in accidents.⁶¹ Other studies found some degree of fatty streaks in the aorta in all persons aged 12–15 years⁶⁵ and fatty streaks in the coronary arteries in approximately 30% of children aged 8–11 years and 69% of persons aged 12–15 years.⁶² The clinical importance of these lesions depends on their anatomical location.

Although epidemiological studies have not directly confirmed the link between the early presence of aortic fatty streaks and the occurrence of clinically consequential atherosclerotic plaques in later life,^{66,67} there is a relationship between the location of fatty streaks in the coronary arteries in children and atherosclerotic lesions in the same site later in life.⁶⁸ In nonblack populations, the extent of involvement of coronary artery with fatty streaks in youth predicts the extent of involvement with raised lesions in older persons.⁶⁹ In an autopsy study of coronary arteries of 565 individuals aged 0–29 years, the progressive transformation of fatty streaks in

Table 1 Effects of vegetarian diets on cardiovascular risk factors

Study design	Populations	CVD risk factor	Key findings	Reference group	References
Systematic review and meta-analysis of cross-sectional studies	71 studies for vegetarian and 19 studies for vegan diet	BMI	-1.49 kg/m ² (95%CI, -1.72 to -1.25) for vegetarians; -1.72 kg/m ² (95%CI, -2.21 to -1.22) for vegans	Various otherwise similar nonvegetarian populations	Dinu et al. (2017) ²⁰
Systematic review and meta-analysis of cross-sectional studies	64 studies for vegetarian and 19 studies for vegan diet	Total cholesterol	-0.72 mmol/L (95%CI, -0.8 to -0.64) for vegetarians; -0.80mmol/L (95%CI, -0.90 to -0.70) for vegans	Various otherwise similar nonvegetarian populations	Dinu et al. (2017) ²⁰
Systematic review and meta-analysis of cross-sectional studies	46 studies for vegetarian and 13 studies for vegan diet	LDL cholesterol	-0.55 mmol/L (95%CI, -0.62 to -0.47) for vegetarians; -0.59 mmol/L (95%CI, -0.77 to -0.40) for vegans	Various otherwise similar nonvegetarian populations	Dinu et al. (2017) ²⁰
Systematic review and meta-analysis of randomized clinical trials	7 trials of intervention with vegan diet and 4 trials of intervention with vegetarian diet	Total cholesterol	-0.36 mmol/L (95%CI, 0.55 to -0.17) with intervention with vegetarian or vegan diet (pooled estimates)	Trial control period	Wang et al. (2015) ¹⁴
Systematic review and meta-analysis of 11 randomized clinical trials	7 trials of intervention with vegan diet and 4 trials of intervention with vegetarian diet	LDL cholesterol	-0.11) with intervention with vegetarian or vegan diet (pooled estimates)	Trial control period	Wang et al. (2015) ¹⁴
Systematic review and meta-analysis of 11	7 trials of intervention with vegan diet and 4 trials of intervention with vegetarian diet	non-HDL cholesterol	-0.30 mmol/L (95%CI, -0.50 to -0.10) with intervention with vegetarian or vegan diet (pooled estimates)	Trial control period	Wang et al. (2015) ¹⁴
Adventist Health Study 2, a cohort study	96 000 participants (7% vegan, 29.2% vegetarian)	Risk of hypertension	RR = 0.86 (95%CI, 0.51–1.45) for vegetarians; RR = 0.53 (95%CI, 0.25–1.11) for vegans	Nonvegetarian Adventists	Orlich et al. (2014) ³⁴
Matched cohort study	4109 Taiwanese nonsmokers	Risk of hypertension	RR = 0.72 (95%CI, 0.55–0.86) for vegetarians	5 omnivores matched to 1 vegetarian by age, sex, and study site	Chuang et al. (2016) ³⁵
A systematic review and meta-analysis of controlled clinical trials and observational studies	7 trials, a total of 311 participants, mean age of 44.5 y; 32 observational studies, a total of 21 604 participants, mean age of 46.6 y	Systolic blood pressure	A reduction in mean systolic blood pressure (-4.8 mm Hg; 95%CI, -6.6 to -3.1) for vegetarian diet	Trial control period	Yokoyama et al. (2014) ¹⁵
A systematic review and meta-analysis of controlled clinical trials and observational studies	7 trials, a total of 311 participants, mean age of 44.5 y; 32 observational studies, a total of 21 604 participants, mean age of 46.6 years	Diastolic blood pressure	Lower mean systolic blood pressure (-6.9 mm Hg; 95%CI, -9.1 to -4.7) for vegetarian diet A reduction in mean diastolic blood pressure (-2.2 mm Hg; 95%CI, -3.5 to -1.0) for vegetarian diet	Various otherwise similar nonvegetarian populations Trial control period	Yokoyama et al. (2014) ¹⁵

(continued)

Table 1 Continued

Study design	Populations	CVD risk factor	Key findings	Reference group	References
Two prospective cohort studies	Adventist Health Study 1 (25 698 participants; approximately 50% vegetarians) and 2 (60 903 participants; 52% vegetarians)	Prevalence of diabetes	Lower mean systolic BP (−4.7 mm Hg; 95%CI, −6.3 to −3.1) for vegetarian diet 1.5–2 times lower	Various otherwise similar nonvegetarian populations Nonvegetarian Adventists	Snowdon and Phillips (1985) ⁴⁶ Tonstad et al. (2009) ⁴⁵
Systematic review and meta-analysis of observational studies	27 studies with 2256 vegetarian and 2192 nonvegetarian participants; 4 studies of 83 vegans and 125 omnivores	Blood glucose levels	−5.08 mg/dL (95%CI, −5.98 to −4.19) for vegetarians; −6.39 mg/dL (95%CI, −12.35 to −0.41) for vegans	Various otherwise similar nonvegetarian populations	Dinu et al. (2017) ²⁰
Review	2 randomized clinical trials of interventions with vegetarian diet in diabetes including 43 and 74 participants with diabetes	Effect on various diabetes markers	A greater reduction in various measures of diabetes, including body weight, fasting plasma glucose, HbA1c, and hypoglycemic medication and greater increase in insulin sensitivity with vegetarian diet	Hypocaloric diet or a diet recommended by American Diabetes Association	Kahleova et al. (2015) ⁴⁷
Cross-sectional study	30 long-term (≥ 5 years) vegetarians and 30 age-matched omnivores	CRP levels	Lower CRP levels in vegetarians (0.77 mg/L; SD, 1.29; $P < 0.01$) than in omnivores (1.30 mg/L; SD, 1.38)	Matched omnivores	Szeto et al. (2004) ⁵⁶
Randomized control trial	46 healthy, hyperlipidemic adults randomized to a diet low in saturated fat and a vegetarian diet high in plant sterols	Effect on CRP levels	Vegetarian diet intervention reduced CRP levels by 28.2% (SE, 10.8%; $P = .02$), whereas control diet reduced CRP by 10% (SE, 8.6%; $P = 0.27$)	Control arm of the clinical trial	Jenkins et al. (2003) ⁵⁷

Abbreviations: BMI, body mass index; BP, blood pressure; CI, confidence interval; CRP, C-reactive protein; CVD, cardiovascular disease; HDL, high-density lipoprotein; LDL, low-density lipoprotein; RR, relative risk; SD, standard deviation; SE, standard error.

children's coronary arteries to a well-advanced fibrous plaque in young adulthood was observed.⁷⁰ In this study, by puberty a small 8%–10% percentage of children had evidence of more advanced lesions, and approximately 30% of the adults in their 20s had well-developed raised lesions with large extracellular lipid cores and thick fibromuscular caps.⁷⁰

Overall, the evidence indicates that the atherosclerotic process starts in the early years. This emphasizes the importance of understanding the factors contributing to variability of its severity among individuals during childhood.

CHILDHOOD ANTECEDENTS OF ADULT ATHEROSCLEROSIS AND CLINICAL CARDIOVASCULAR DISEASE

The only direct evidence linking cardiometabolic risk factors other than BMI in youth⁷¹ with overt clinical disease in adulthood comes from genetic disorders related to high cholesterol. In homozygous familial hypercholesterolemia, a genetic disease whereby LDL-C clearance is impaired, LDL-C levels exceed 20.68 mmol/L already in infants, CVD events begin in the first decade of life, and life span is reduced.⁷² In heterozygous hypercholesterolemia, in which LDL-C levels usually exceed 5.17 mmol/L and TC levels exceed 6.5 mmol/L beginning in infancy, 50% of men and 25% of women experience clinical coronary events by the age of 50 years.⁷² Furthermore, in familial hypertriglyceridemia, another genetic disorder resulting in excessive triglyceride (TG) levels, childhood TG levels independently predict CVD in the 4th–5th decade of life.⁷³

The substantial genetic component of CVD risk can also be seen among children of patients suffering from premature myocardial infarction (<55 y), who have higher levels of TC, LDL-C, and TGs and lower levels of HDL-C than controls.⁷⁴ Another piece of evidence linking childhood blood lipids with CVD risk in adulthood in the context of genetic predisposition stems from Mendelian randomization studies. Meta-analysis of such studies found a 54.5% (95%CI, 48.8%–59.5%) reduction in the risk of IHD per millimole-per-liter reduction in LDL-C (effect size 3-fold greater than that achieved via treatment with statins in later life) due to genetic polymorphism, and thus relating to lower low-density lipoprotein cholesterol lifetime exposure.⁷⁵

Most evidence suggesting that childhood cardiometabolic physiology affects adult arterial pathology is indirect and comes from autopsy and imaging studies linking childhood risk factors with atherosclerosis and its surrogate markers. The extent to which the artery surface is involved in individual children with lesions varies; however, it is influenced by the same classical

risk factors that predict adult coronary heart disease.^{62,76} The Bogalusa Heart Study has demonstrated a strong association of BMI, SBP, DBP, and serum concentrations of TC, TGs, LDL-C, and HDL-C with vascular lesions in children and young adults on autopsy.⁷⁷

The PDAY autopsy study showed strong relationships between atherosclerotic severity and extent with age, non-HDL-C, HDL-C, hypertension (determined by renal artery thickness), tobacco use (thiocyanate concentration), diabetes mellitus (glycohemoglobin), and (in men) obesity. It also showed that a 30 mg/dL incremental increase in non-HDL-C was equivalent to 2 years of “vascular aging.”⁷⁸ The severity and extent of the lesions were positively associated with age and increased in association with the number of risk factors. At the same time, an absence of risk factors was associated with a virtual absence of advanced atherosclerotic lesions, even in the oldest individuals in the study.

Analyses from 4 longitudinal cohorts (Cardiovascular Risk in Young Finns Study, Childhood Determinants of Adult Health Study, Bogalusa Heart Study, and Muscatine Study) showed that risk factors measured at age 9 years or after (TC, TGs, blood pressure, and BMI) were predictive of elevated carotid IMT in adulthood,⁷⁹ which is recognized as a predictive measure of clinical coronary events in middle-aged and elderly populations.⁸⁰ Similarly, SBP, DBP, TC, LDL-C, HDL-C, and smoking status were linked to IMT of the femoral artery, a surrogate measure of coronary and peripheral atherosclerosis, in asymptomatic young individuals in the Bogalusa Heart Study.⁸¹

Additionally, in the Cardiovascular Risk in Young Finns Study, childhood LDL-C (≥ 80 th percentile), elevated blood pressure, skinfold thickness, low HDL-C (≤ 20 th percentile), and smoking were inversely associated with artery elasticity in adulthood,⁸² a marker of pathophysiological changes in the arteries relevant to the development of atherosclerosis later in life.⁸³ Increased body size, increased blood pressure, and decreased HDL-C were associated with coronary artery calcification in young adults in the Muscatine Study.⁸⁴

Other studies have examined the relationship of isolated childhood risk factors to various measures of atherosclerosis and CVD risk. Dietary fat quality reflected in the serum cholesterol ester fraction in childhood was associated with carotid IMT in adult women.⁸⁵ Children with hypercholesterolemia and diabetes showed increased IMTs compared with healthy controls,⁸⁶ and cumulative exposure to hyperlipidemia in young adulthood increased subsequent risk of coronary heart disease in a dose-dependent fashion.⁸⁷ Other, nondiet-related risk factors such as smoking⁸⁸ and exercise⁸⁹ in childhood were also associated with

adult atherosclerosis in a pattern similar to that in adulthood.

Collectively, all of these studies indicate that modifiable phenotypic traits in childhood, including diet and body fatness, are associated with the early emergence of atherosclerotic progression to calcified plaque, manifesting as overt CVD in adult life. Children with fewer cardiometabolic risk factors show lower prevalence of atherosclerotic risk in later life. Dietary-influenced risk factors in childhood and their relation to atherosclerosis are summarized in Table 2^{77-79,81,82,84,86}.

TRACKING OF CHILDHOOD CARDIOVASCULAR DISEASE RISK FACTORS AND THEIR DETERMINANTS INTO ADULTHOOD

Cardiometabolic risk factors in childhood deserve attention not only because of their association with atherosclerosis in adulthood but also because there is evidence that they track (ie, persist) into adult life, therefore generating a cumulative impact on the process of disease. Evidence for tracking is strongest for obesity, with childhood BMI levels predictive of adult obesity.⁹⁰ A recent systematic review of 13 prospective or retrospective longitudinal studies published after 2001 showed that the risk of an overweight child becoming an overweight adult is at least twice as high compared with normal-weight children, and it is even higher for obese children.⁹¹ This could be due to both the direct tracking of body composition and the tracking of obesity-related behaviors—such as physical inactivity and unhealthy diets—between childhood and adulthood.⁹²

Correlation coefficients for cholesterol tracking are in the range of 0.4 and have been found consistently in numerous studies examining children as young as 5–10 years of age and their lipid levels 20–30 years later.⁷² In the Muscatine Study, 75% of children aged 5–10 years who had TC concentrations >90th percentile at baseline had TC concentrations >200 mg/dL in their early 20s.^{93,94} In the Bogalusa Heart Study, approximately 50% of those children who had TC levels or LDL-C levels above the 75th percentile at baseline had elevated levels 12 years later.⁹⁵ In the same study, adverse glucose levels in childhood not only persisted into adulthood but also predicted adult prediabetes and type 2 diabetes.⁹⁶

In a retrospective cohort study of 1058 normoglycemic, 37 prediabetic, and 25 type 2 diabetic adults aged 19–39 years followed on average for 17 years since childhood, at least 50% of the individuals who ranked in the top childhood quintile for glucose, insulin, and homeostatic model assessment (HOMA) insulin

resistance had measurements above the 60th percentile in adulthood.⁹⁶

Elevated blood pressure in youth predicts adult hypertension, and a systematic review and meta-analysis of 50 cohort studies reported degree of tracking with correlation coefficient at 0.38 for SBP and 0.28 for DBP.⁹⁷

Tracking of CRP levels was observed between childhood and adulthood in a cohort of 1617 individuals aged 3–18 years at baseline and reexamined at 24–39 years of age. The age- and sex-specific correlations were the highest in the group aged 18 years at baseline ($r = 0.47$ in females, $r = 0.32$ in males).⁹⁸

Of particular relevance to this review, diet itself, one of the strongest correlates of cardiometabolic risk, also tracks from childhood into adulthood. The Cardiovascular Risk in Young Finns Study, a prospective cohort study with a 21-year follow-up, found some level of tracking of dietary patterns.⁹⁹ Similarly, a review of studies published between 2003 and 2013 reported moderate level of tracking for a range of eating behaviors (eg, food preferences, dietary variety, dietary intake, eating habits) measured before 10 years of age and reassessed in adulthood.¹⁰⁰

IMPORTANCE OF PRIMORDIAL PREVENTION

As discussed at the outset of this review, the considerable gains made in reducing CVD mortality rates since the 1970s through risk factor reductions and better treatment of the disease are increasingly challenged by adverse trends in obesity and diabetes.^{101,102} Better treatment strategies are not expected to offset these adverse trends^{7,103} and would increase exponentially medical costs given the aging of most populations. Moreover, treatment strategies reduce but do not eliminate the risk of CVD.^{104,105}

Therefore, risk factor reduction strategies are critical for reducing the CVD burden. Numerous modeling studies have estimated that population-wide risk factor reductions can bring substantial decreases of CVD burden,^{8,106-108} even taking into account current trends of obesity and diabetes. It has also been shown that mortality trends respond very rapidly to changes in risk factors at the population level.¹⁰⁹

Given compelling evidence that the atherosclerotic process starts in childhood and is linked to well-defined, modifiable risk factors that track into adulthood, there is increasing recognition that primordial prevention (ie, avoiding the development of risk factors before the disease onset) should be embraced as a major component of global CVD prevention policies.¹¹⁰ This is an approach through which favorable patterns for all lifestyle (eg, physical activity, avoidance of smoking) and all

Table 2 Dietary-influenced risk factors in childhood and their relation to atherosclerosis

Study	Population	Findings	Locations of the lesions measured	Additional information	References
Bogalusa Heart Study	Autopsies on 204 young persons aged 2–39 y	A strong association of BMI, SBP, DBP, and serum concentrations of total cholesterol, TG, LDL-C, and HDL-C with vascular lesions in children and young adults on autopsy	Coronary arteries, aorta	The association between less advanced lesions (fatty streaks) and more advanced ones (fibrous plaques) was much stronger in the coronary arteries than in the aorta	Berenson et al. (1998) ⁷⁷
Pathobiological Determinants of Atherosclerosis in Youth study	> 3000 autopsies of persons aged 15–34 y	Strong relationships between atherosclerotic severity/extent with age, non-HDL-C, HDL-C, hypertension, tobacco use, diabetes mellitus, and (in men) obesity on autopsy	Left anterior descending coronary artery, right coronary artery, and abdominal aorta	Severity and extent of lesions positively associated with age and with number of risk factors	McMahan et al. (2006) ⁷⁸
4 longitudinal cohorts (Cardiovascular Risk in Young Finns Study, Childhood Determinants of Adult Health study, Bogalusa Heart Study, and Muscatine Study)	4380 members of 4 prospective cohorts with cardiovascular risk factor data from childhood (aged 3–18 y) and IMT in adulthood (aged 20–45 y)	Risk factors at age > 9 y (total cholesterol, TG, blood pressure, and BMI) were predictive of elevated carotid IMT in adulthood	Carotid IMT	The associations with risk factors measured at age 3 y and 6 y were weaker and nonsignificant	Juonala et al. (2010) ⁷⁹
Bogalusa Heart Study	1080 black and white individuals (aged 24–43 y; 71% white, 43% male); individuals in the top (n = 54) vs bottom fifth (n = 54) percentiles distribution of femoral IMT were compared for traditional cardiovascular risk factors profile	SBP, DBP, total cholesterol, LDL-C, HDL-C, and smoking status were linked to IMT of the femoral artery	Femoral artery IMT		Paul et al. (2005) ⁸¹
Cardiovascular Risk in Young Finns Study	2255 healthy white adults aged 24–39 y who had risk factor data available since childhood	Childhood LDL-C (\geq 80th percentile), elevated blood pressure, skinfold thickness, low HDL-C (\leq 20th percentile), and smoking were inversely associated with artery elasticity in adulthood	Carotid artery elasticity comprising carotid artery compliance, Young's elastic modulus, and stiffness index	Associations remained highly significant after adjustment for the number of risk factors identified in adulthood	Juonala et al. (2005) ⁸²

(continued)

Table 2 Continued

Study	Population	Findings	Locations of the lesions measured	Additional information	References
Muscatine Study	384 individuals (197 men, 187 women) who had coronary risk factors measured in childhood (mean age, 15 y) and twice during young adult life (mean ages, 27 and 33 y)	Increased body size, increased blood pressure, and decreased HDL-C were associated with coronary artery calcification in young adults			Mahoney et al. (1996) ⁸⁴
Cross-sectional study	88 children (aged 11 ± 2 y)	Children with hypercholesterolemia and diabetes showed increased IMTs compared with healthy controls	Aortic and carotid IMT		Järvisalo et al. (2001) ⁸⁶

Abbreviations: BMI, body mass index; DBP, diastolic blood pressure; HDL-C, high-density lipoprotein cholesterol; IMT, intima media thickness; LDL-C, low-density lipoprotein cholesterol; SBP, systolic blood pressure; TG, triglycerides.

major lifestyle-related risk factors (eg, cholesterol, hypertension, glucose levels) can be promoted from conception throughout childhood. Those patterns can then potentially be maintained into adulthood.¹¹¹

The concept of primordial prevention is strongly backed by data. Two prospective studies from the late 1960s and 1970s—the Chicago Heart Association Detection Project in Industry Study¹¹² (n = 8816) and the Multiple Risk Factor Intervention Trial Study¹¹³ (n = 12 866)—showed that favorable levels of all readily-measured major CVD risk factors in young adulthood (TC < 5.17 mmol/L; SBP ≤ 120 mm Hg; DBP ≤ 80 mm Hg; BMI < 25.0 kg/m²) lead to substantially reduced CVD mortality rates (76%–89% for men and 60%–67% lower in women) and sizable increases in life expectancy (8–12 y greater).

So far, CVD prevention strategies targeting early life have primarily been focused on tackling childhood obesity, although success rates are poor.¹¹⁴ Interest has also focused on factors like birth weight²³ and early catch-up growth,²⁶ although some of these traits are difficult to target through interventions because of the need to change maternal physiology. It is known, however, from the Cardiovascular Risk in Young Finns Study, Childhood Determinants of Adult Health Study, Bogalusa Heart Study, and Muscatine Study, that other classical CVD risk factors relate to adult atherosclerosis independently from BMI⁷⁹ and would therefore benefit from additional interventions. For example, recent evidence from the United States suggests that up to one third of prepubertal children with normal weight have abnormal lipid levels¹¹⁵ and that the prevalence of hypertension in the pediatric population has been increasing.¹¹⁶

Therefore, interventions effectively targeting all classical risk factors in young people could potentially play a key role in preventive strategies. Linking the 2 components of this review—that adults consuming plant-based diets have lower CVD risk and that cardiometabolic risk tracks from childhood into adulthood—a new testable hypothesis is now proposed: namely, that a plant-based diet in childhood could promote cardiometabolic health in adults, thereby reducing CVD and promoting longevity and health.

PLANT-BASED DIETS IN CHILDHOOD AS A MEANS TO PROMOTE CARDIOMETABOLIC HEALTH IN ADULTHOOD

Diet in children appears to be one of the strongest determinants of the CVD risk factors.¹¹⁷ Additionally, childhood diets show some degree of tracking into adulthood^{99,100} and associate with adulthood cardiovascular risk factors, vascular markers of subclinical atherosclerosis, and CVD risk.²⁷ The few available studies in this




Figure 2 Potential direct and indirect mechanisms through which plant-based diets in children could benefit cardiometabolic health in adults.

area suggest that healthy childhood dietary patterns are associated with lower adulthood CVD risk.^{27,118} Intakes of plant foods (vegetables, fruits, and fiber) and polyunsaturated fatty acids have shown protective effects.^{27,118}

Vegan and vegetarian children have lower rates of overweight and obesity.²⁸ Preliminary evidence suggests that they have lower cholesterol levels^{119,120} and higher antioxidant status in the blood.¹²¹ They consume more fruits and vegetables than their omnivore counterparts.^{122,123} Moreover, a recent trial showed that an intervention with low-fat vegan diet was more effective than the American Heart Association–recommended diet at reducing CVD risk factors in obese and hypercholesterolemic children aged 9–18 years old. Children assigned to the vegan intervention had more significant ($P < 0.05$) reductions in CVD risk factors from baseline—BMI Z-score (−0.14), systolic SBP (−6.43 mm Hg), TC (−22.5 mg/dL), LDL-C (−13.14 mg/dL), high sensitivity CRP (−2.09 mg/L), insulin (−5.42uU/ml), myeloperoxidase (−75.34 pmol/L), midarm circumference (−2.02 cm), weight (−3.05 kg)—whereas in the American Heart Association group significant ($P < 0.05$) were noted only for the last 3 risk factors (−69.23 pmol/L, −1.55 cm, −1.14 kg, respectively) and waist circumference (−2.96 cm).¹²⁴

Therefore, vegetarian and vegan children might have a better CVD risk profile than omnivore children, and if the diet and risk profile tracks into adulthood, plant-based diets in pediatric populations, through their

effects on blood biochemistry, other cardiovascular risk factors, and the establishment of healthy eating patterns, could offer an effective strategy of CVD primordial prevention. This would present a novel opportunity to promote a healthy childhood trajectory toward cardiovascular health (see Figure 2).

At the same time, data on the safety of vegetarian diets in childhood are sparse and suggest increased risk of nutrient deficiencies, especially in vegan children, including some deficiencies that can differentially affect CVD risk, such as deficiencies in vitamin B12, vitamin D, n-3 essential fatty acids, and iron.^{122,125} Therefore 2 issues are of importance. First, further research is required to comprehensively assess the safety, along with the CVD-protective potential, of vegetarian and vegan diets in childhood. Second, preventive strategies should potentially consider not only promoting vegetarian diets per se but also increasing the percentage of plant foods within omnivore diets in children.

CONCLUSION

The population-wide reduction of CVD risk factors seems currently to be the only feasible strategy to combat the CVD epidemic. More attention should be given to pediatric populations in this context to take advantage of primordial prevention. Atherosclerosis starts in early life and progresses in relation to the same classical risk factors in children that influence the course of the

disease in adults. Moreover, these risk factors, along with dietary habits, track into adulthood. Children with fewer cardiometabolic risk factors show lower prevalence of atherosclerotic risk in later life. Plant-based diets have been shown to substantially reduce CVD risk factors, morbidity, and mortality in adults and also offer planetary sustainability benefits. If applied in children, they could potentially offer cardiometabolic health benefits through reduction of CVD risk beginning in early life, a strategy that has been shown to bring the most favorable CVD mortality reductions and increases in life expectancy. At the same time, little is known about the safety and potential cardiometabolic benefits of these diets applied from childhood, and further research in this area is warranted.

Acknowledgments

Author contributions. M.D. contributed significantly to the work's conception and design, wrote and revised the manuscript, and managed references. J.S. critically reviewed and contributed to the writing of the manuscript. M.F. participated in the critical revision of the article. J.C.K.W. contributed significantly to the work's conception and design and participated in the writing and critical revision of the article.

Funding. No external funding supported this review.

Declaration of interest. The authors have no relevant interests to declare.

REFERENCES

- World Health Organization. *Global Status Report on Noncommunicable Diseases*. http://www.who.int/nmh/publications/ncd_report2010/en/. Accessed April 6, 2017.
- World Health Organization. *Disability-Adjusted Life Years (DALYs)*. http://www.who.int/gho/mortality_burden_disease/daly_rates/text/en/. Accessed April 6, 2017.
- Levi F, Chatenoud L, Bertuccio P, et al. Mortality from cardiovascular and cerebrovascular diseases in Europe and other areas of the world: an update. *Eur J Cardiovasc Prev Rehabil*. 2009;16:333–350.
- Allender S, Scarborough P, O'Flaherty M, et al. Patterns of coronary heart disease mortality over the 20th century in England and Wales: possible plateaus in the rate of decline. *BMC Public Health*. 2008;8:148.
- Ford ES, Capewell S. Coronary heart disease mortality among young adults in the US from 1980 through 2002: concealed leveling of mortality rates. *J Am Coll Cardiol*. 2007;50:2128–2132.
- O'Flaherty M, Allender S, Taylor R, et al. The decline in coronary heart disease mortality is slowing in young adults (Australia 1976–2006): a time trend analysis. *Int J Cardiol*. 2012;158:193–198.
- Emberson J, Whincup P, Morris R, et al. Evaluating the impact of population and high-risk strategies for the primary prevention of cardiovascular disease. *Eur Heart J*. 2004;25:484–491.
- O'Flaherty M, Bandosz P, Critchley J, et al. Exploring potential mortality reductions in 9 European countries by improving diet and lifestyle: a modelling approach. *Int J Cardiol*. 2012;158:207–286–291.
- Charakida M, Deanfield JE, Halcox JJP. Childhood origins of arterial disease. *Curr Opin Pediatr*. 2007;19:538–545.
- Mozaffarian D. Dietary and policy priorities for cardiovascular disease, diabetes, and obesity. *Circulation*. 2016;133:187–225.
- Sabaté J, Soret S. Sustainability of plant-based diets: back to the future. *Am J Clin Nutr*. 2014;100(suppl 1):476S–482S.
- Ginsberg, C. *The Market for Vegetarian Foods*. <http://www.vrg.org/nutshell/market.htm#market>. Accessed April 11, 2017.
- Thompson, H. *Vegetarian and Vegan Trend Soars as Consumers Reduce Meat Intake*. <http://www.bighospitality.co.uk/Trends-Reports/Vegetarian-and-vegan-trend-soars-as-consumers-reduce-meat-intake>. Accessed April 11, 2017.
- Wang F, Zheng J, Yang B, et al. Effects of vegetarian diets on blood lipids: a systematic review and meta-analysis of randomized controlled trials. *J Am Heart Assoc*. 2015;4:e002408.
- Yokoyama Y, Nishimura K, Barnard ND, et al. Vegetarian diets and blood pressure: a meta-analysis. *JAMA Intern Med*. 2014;174:577–587.
- Crowe FL, Appleby PN, Travis RC, et al. Risk of hospitalization or death from ischemic heart disease among British vegetarians and nonvegetarians: results from the EPIC-Oxford cohort study. *Am J Clin Nutr*. 2013;97:597–603.
- Key TJ, Fraser GE, Thorogood M, et al. Mortality in vegetarians and nonvegetarians: detailed findings from a collaborative analysis of 5 prospective studies. *Am J Clin Nutr*. 1999;70:516S–524S.
- Fraser GE. Vegetarian diets: what do we know of their effects on common chronic diseases? *Am J Clin Nutr*. 2009;89:1607S–1612S.
- Appleby PN, Crowe FL, Bradbury KE, et al. Mortality in vegetarians and comparable nonvegetarians in the United Kingdom. *Am J Clin Nutr*. 2016;103:218–230.
- Dinu M, Abbate R, Gensini GF, et al. Vegetarian, vegan diets and multiple health outcomes: a systematic review with meta-analysis of observational studies. *Crit Rev Food Sci Nutr*. 2017;57:3640–3649.
- Holman RL, McGill HC, Strong JP, et al. The natural history of atherosclerosis. *Am J Pathol*. 1958;34:209–235.
- Palatianou ME, Simos YV, Andronikou SK, et al. Long-term metabolic effects of high birth weight: a critical review of the literature. *Horm Metab Res*. 2014;46:911–920.
- Wang S-F, Shu L, Sheng J, et al. Birth weight and risk of coronary heart disease in adults: a meta-analysis of prospective cohort studies. *J Dev Orig Health Dis*. 2014;5:408–419.
- Lloyd LJ, Langley-Evans SC, McMullen S. Childhood obesity and adult cardiovascular disease risk: a systematic review. *Int J Obes (London)*. 2010;34:18–28.
- Park MH, Falconer C, Viner RM, et al. The impact of childhood obesity on morbidity and mortality in adulthood: a systematic review. *Obes Rev*. 2012;13:985–1000.
- Jain V, Singhal A. Catch up growth in low birth weight infants: striking a healthy balance. *Rev Endocr Metab Disord*. 2012;13:141–147.
- Kaikkonen JE, Mikkilä V, Raitakari OT. Role of childhood food patterns on adult cardiovascular disease risk. *Curr Atheroscler Rep*. 2014;16:443.
- Sabaté J, Wien M. Vegetarian diets and childhood obesity prevention. *Am J Clin Nutr*. 2010;91:1525S–1529S.
- Klein S, Burke LE, Bray GA, et al. Clinical implications of obesity with specific focus on cardiovascular disease: a statement for professionals from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism: endorsed by the American College of Cardiology Foundation. *Circulation*. 2004;110:2952–2967.
- Bradbury KE, Crowe FL, Appleby PN, et al. Serum concentrations of cholesterol, apolipoprotein A-I and apolipoprotein B in a total of 1694 meat-eaters, fish-eaters, vegetarians and vegans. *Eur J Clin Nutr*. 2014;68:178–183.
- Schwingshackl L, Hoffmann G. Comparison of effects of long-term low-fat vs high-fat diets on blood lipid levels in overweight or obese patients: a systematic review and meta-analysis. *J Acad Nutr Diet*. 2013;113:1640–1661.
- Li D. Chemistry behind vegetarianism. *J Agric Food Chem*. 2011;59:777–784.
- Craig WJ. Nutrition concerns and health effects of vegetarian diets. *Nutr Clin Pract*. 2010;25:613–620.
- Orlich MJ, Fraser GE. Vegetarian diets in the Adventist Health Study 2: a review of initial published findings. *Am J Clin Nutr*. 2014;100(suppl 1):353S–358S.
- Chuang S-Y, Chiu THT, Lee C-Y, et al. Vegetarian diet reduces the risk of hypertension independent of abdominal obesity and inflammation. A prospective study. *J Hypertens*. 2016;34:2164–2171.
- Sobiecki JG, Appleby PN, Bradbury KE, et al. High compliance with dietary recommendations in a cohort of meat eaters, fish eaters, vegetarians, and vegans: results from the European Prospective Investigation into Cancer and Nutrition-Oxford study. *Nutr Res*. 2016;36:464–477.
- Evans CEL, Greenwood DC, Threapleton DE, et al. Effects of dietary fibre type on blood pressure: a systematic review and meta-analysis of randomized controlled trials of healthy individuals. *J Hypertens*. 2015;33:897–911.
- He J, Gu D, Wu X, et al. Effect of soybean protein on blood pressure: a randomized, controlled trial. *Ann Intern Med*. 2005;143:1–9.
- Haddad EH, Tanzman JS. What do vegetarians in the United States eat? *Am J Clin Nutr*. 2003;78(3 suppl):626S–632S.
- Aburto NJ, Hanson S, Gutierrez H, et al. Effect of increased potassium intake on cardiovascular risk factors and disease: systematic review and meta-analyses. *BMJ*. 2013;346:f1378.

41. Galan P, Vergnaud A-C, Tzoulaki I, et al. Low total and nonheme iron intakes are associated with a greater risk of hypertension. *J Nutr*. 2010;140:75–80.
42. Sciarrone SE, Strahan MT, Bellin LJ, et al. Biochemical and neurohormonal responses to the introduction of a lacto-ovo-vegetarian diet. *J Hypertens*. 1993;11:849–860.
43. Ernst E, Pietsch L, Matrai A, et al. Blood rheology in vegetarians. *Br J Nutr*. 1986;56:555–560.
44. Ettehad D, Emdin CA, Kiran A, et al. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet*. 2016;387:957–967.
45. Tonstad S, Butler T, Yan R, et al. Type of vegetarian diet, body weight, and prevalence of type 2 diabetes. *Diabetes Care*. 2009;32:791–796.
46. Snowdon DA, Phillips RL. Does a vegetarian diet reduce the occurrence of diabetes? *Am J Public Health*. 1985;75:507–512.
47. Kahleova H, Pelikanova T. Vegetarian diets in the prevention and treatment of type 2 diabetes. *J Am Coll Nutr*. 2015;34:448–458.
48. Kahleova H, Matoulek M, Malinska H, et al. Vegetarian diet improves insulin resistance and oxidative stress markers more than conventional diet in subjects with type 2 diabetes. *Diabet Med*. 2011;28:549–559.
49. Barnard ND, Cohen J, Jenkins DJA, et al. A low-fat vegan diet improves glycemic control and cardiovascular risk factors in a randomized clinical trial in individuals with type 2 diabetes. *Diabetes Care*. 2006;29:1777–1783.
50. Fung TT, Schulze M, Manson JE, et al. Dietary patterns, meat intake, and the risk of type 2 diabetes in women. *Arch Intern Med*. 2004;164:2235–2240.
51. Sievenpiper JL, Kendall CWC, Esfahani A, et al. Effect of non-oil-seed pulses on glycaemic control: a systematic review and meta-analysis of randomised controlled experimental trials in people with and without diabetes. *ResearchGate*. 2009;52:1479–1495.
52. Aune D, Norat T, Romundstad P, et al. Whole grain and refined grain consumption and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies. *Eur J Epidemiol*. 2013;28:845–858.
53. Cooper AJ, Forouhi NG, Ye Z, et al. Fruit and vegetable intake and type 2 diabetes: EPIC-InterAct prospective study and meta-analysis. *Eur J Clin Nutr*. 2012;66:1082–1092.
54. Pan A, Sun Q, Bernstein AM, et al. Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *Am J Clin Nutr*. 2011;94:1088–1096.
55. Collaboration TERF. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies. *Lancet*. 2010;375:2215–2222.
56. Szeto YT, Kwok TCY, Benzie IFF. Effects of a long-term vegetarian diet on biomarkers of antioxidant status and cardiovascular disease risk. *Nutrition*. 2004;20:863–866.
57. Jenkins DJA, Kendall CWC, Marchie A, et al. Effects of a dietary portfolio of cholesterol-lowering foods vs lovastatin on serum lipids and C-reactive protein. *JAMA*. 2003;290:502–510.
58. Acosta-Navarro J, Antoniazzi L, Oki AM, et al. Reduced subclinical carotid vascular disease and arterial stiffness in vegetarian men: the CARVOS study. *Int J Cardiol*. 2017;230:562–566.
59. Yang S-Y, Zhang H-J, Sun S-Y, et al. Relationship of carotid intima-media thickness and duration of vegetarian diet in Chinese male vegetarians. *Nutr Metab (London)*. 2011;8:63.
60. Su T-C, Jeng J-S, Wang J-D, et al. Homocysteine, circulating vascular cell adhesion molecule and carotid atherosclerosis in postmenopausal vegetarian women and omnivores. *Atherosclerosis*. 2006;184:356–362.
61. McGill HC, McMahan CA, Herderick EE, et al. Origin of atherosclerosis in childhood and adolescence. *Am J Clin Nutr*. 2000;72:1307s–1315s.
62. Stary HC. Lipid and macrophage accumulations in arteries of children and the development of atherosclerosis. *Am J Clin Nutr*. 2000;72:1297s–1306s.
63. Enos WF, Holmes RH, Beyer J. Coronary disease among United States soldiers killed in action in Korea preliminary report. *J Am Med Assoc*. 1953;152:1090–1093.
64. McNamara JJ, Molot MA, Stremple JF, et al. Coronary artery disease in combat casualties in Vietnam. *JAMA*. 1971;216:1185–1187.
65. Stary HCYZ. The natural history of atherosclerosis in the aorta in the first forty years of life. In Fuster V, ed. *Syndromes of Atherosclerosis: Correlations of Clinical Imaging and Pathology*. New York: Futura Publishing; 1996:225–238.
66. Stehbens WE. The epidemiological relationship of hypercholesterolemia, hypertension, diabetes mellitus and obesity to coronary heart disease and atherogenesis. *J Clin Epidemiol*. 1990;43:733–741.
67. Newman TB, Garber AM, Holtzman NA, et al. Problems with the report of the expert panel on blood cholesterol levels in children and adolescents. *Arch Pediatr Adolesc Med*. 1995;149:241–247.
68. Montenegro MR, Eggen DA. Topography of atherosclerosis in the coronary arteries. *Lab Invest*. 1968;18:586–593.
69. Smith EB. The influence of age and atherosclerosis on the chemistry of aortic intima 2 collagen and mucopolysaccharides. *J Atheroscler Res*. 1965;5:241–248.
70. Stary HC. Evolution and progression of atherosclerotic lesions in coronary arteries of children and young adults. *Arterioscler Dallas Tex*. 1989;9(1 suppl):119–32.
71. Twig G, Yaniv G, Levine H, et al. Body-mass index in 2.3 million adolescents and cardiovascular death in adulthood. *N Engl J Med*. 2016;374:2430–2440.
72. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents, National Heart, Lung, and Blood Institute. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents: summary report. *Pediatrics*. 2011;128(suppl 5):S213–S256.
73. Morrison JA, Glueck CJ, Wang P. Childhood risk factors predict cardiovascular disease, impaired fasting glucose plus type 2 diabetes mellitus, and high blood pressure 26 years later at mean age 38: The Princeton LRC follow-up study. *Metabolism*. 2012;61:531–541.
74. Kelishadi R, Zadeegan NS, Naderi GA, et al. Atherosclerosis risk factors in children and adolescents with or without family history of premature coronary artery disease. *Med Sci Monit*. 2002;8:CR425–42CR429.
75. Ference BA, Yoo W, Alesh I, et al. Effect of long-term exposure to lower low-density lipoprotein cholesterol beginning early in life on the risk of coronary heart disease. A Mendelian randomization analysis. *J Am Coll Cardiol*. 2012;60:2631–2639.
76. Newman WP, Freedman DS, Voors AW, et al. Relation of serum lipoprotein levels and systolic blood pressure to early atherosclerosis. The Bogalusa Heart Study. *N Engl J Med*. 1986;314:138–144.
77. Berenson GS, Srinivasan SR, Bao W, et al. Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. *N Engl J Med*. 1998;338:1650–1656.
78. McMahan CA, Gidding SS, Malcom GT, et al. Pathobiological determinants of atherosclerosis in youth risk scores are associated with early and advanced atherosclerosis. *Pediatrics*. 2006;118:1447–1455.
79. Juonala M, Magnussen CG, Venn A, et al. Influence of age on associations between childhood risk factors and carotid intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study, the Childhood Determinants of Adult Health Study, the Bogalusa Heart Study, and the Muscatine Study for the International Childhood Cardiovascular Cohort (i3C) Consortium. *Circulation*. 2010;122:2514–2520.
80. Hodis HN, Mack WJ, LaBree L, et al. The role of carotid arterial intima-media thickness in predicting clinical coronary events. *Ann Intern Med*. 1998;128:262–269.
81. Paul TK, Srinivasan SR, Wei C, et al. Cardiovascular risk profile of asymptomatic healthy young adults with increased femoral artery intima-media thickness: the Bogalusa Heart Study. *Am J Med Sci*. 2005;330:105–110.
82. Juonala M, Järvisalo MJ, Mäki-Torkko N, et al. Risk factors identified in childhood and decreased carotid artery elasticity in adulthood. *Circulation*. 2005;112:1486–1493.
83. Oliver JJ, Webb DJ. Noninvasive assessment of arterial stiffness and risk of atherosclerotic events. *Arterioscler Thromb Vasc Biol*. 2003;23:554–566.
84. Mahoney LT, Burns TL, Stanford W, et al. Coronary risk factors measured in childhood and young adult life are associated with coronary artery calcification in young adults: the Muscatine Study. *J Am Coll Cardiol*. 1996;27:277–284.
85. Kaikkonen JE, Jula A, Mikkilä V, et al. Childhood serum fatty acid quality is associated with adult carotid artery intima media thickness in women but not in men. *J Nutr*. 2013;143:682–689.
86. Järvisalo MJ, Jartti L, Näntö-Salonen K, et al. Increased aortic intima-media thickness. *Circulation*. 2001;104:2943–2947.
87. Navar-Boggan AM, Peterson ED, D'Agostino RB, et al. Hyperlipidemia in early adulthood increases long-term risk of coronary heart disease. *Circulation*. 2015;131:451–458.
88. Priest JR, Nead KT, Wehner MR, et al. Self-reported history of childhood smoking is associated with an increased risk for peripheral arterial disease independent of lifetime smoking burden. *PLoS One*. 2014;9:e88972.
89. Ried-Larsen M, Grantved A, Kristensen PL, et al. Moderate-and-vigorous physical activity from adolescence to adulthood and subclinical atherosclerosis in adulthood: prospective observations from the European Youth Heart Study. *Br J Sports Med*. 2015;49:107–112.
90. Srinivasan SR, Bao W, Wattigney WA, et al. Adolescent overweight is associated with adult overweight and related multiple cardiovascular risk factors: the Bogalusa Heart Study. *Metabolism*. 1996;45:235–240.
91. Singh AS, Mulder C, Twisk JWR, et al. Tracking of childhood overweight into adulthood: a systematic review of the literature. *Obes Rev*. 2008;9:474–488.
92. Craigie AM, Lake AA, Kelly SA, et al. Tracking of obesity-related behaviours from childhood to adulthood: A systematic review. *Maturitas*. 2011;70:266–284.
93. Lauer RM, Lee J, Clarke WR. Factors affecting the relationship between childhood and adult cholesterol levels: the Muscatine Study. *Pediatrics*. 1988;82:309–318.
94. Lauer RM, Clarke WR. Use of cholesterol measurements in childhood for the prediction of adult hypercholesterolemia. The Muscatine Study. *JAMA*. 1990;264:3034–3038.
95. Webber LS, Srinivasan SR, Wattigney WA, et al. Tracking of serum lipids and lipoproteins from childhood to adulthood the Bogalusa Heart Study. *Am J Epidemiol*. 1991;133:884–899.

96. Nguyen QM, Srinivasan SR, Xu J-H, et al. Utility of childhood glucose homeostasis variables in predicting adult diabetes and related cardiometabolic risk factors. *Diabetes Care*. 2010;33:670–675.
97. Chen X, Wang Y. Tracking of blood pressure from childhood to adulthood: a systematic review and meta-regression analysis. *Circulation*. 2008;117:3171–3180.
98. Juonala M, Viikari JSA, Rönkä T, et al. Childhood C-reactive protein in predicting CRP and carotid intima-media thickness in adulthood. *Arterioscler Thromb Vasc Biol*. 2006;26:1883–1888.
99. Mikkilä V, Räsänen L, Raitakari OT, et al. Consistent dietary patterns identified from childhood to adulthood: the cardiovascular risk in Young Finns Study. *Br J Nutr*. 2005;93:923–931.
100. Nicklaus S, Remy E. Early origins of overeating: tracking between early food habits and later eating patterns. *Curr Obes Rep*. 2013;2:179–184.
101. Fuster V, Kelly BB. *Epidemiology of cardiovascular disease*. In: Promoting Cardiovascular Health in the Developing World: A Critical Challenge to Achieve Global Health. Washington, DC: National Academies Press; 2010. <https://www.ncbi.nlm.nih.gov/books/NBK45688>. Accessed April 8, 2017.
102. Capewell S, Buchan I. Why have sustained increases in obesity and type 2 diabetes not offset declines in cardiovascular mortality over recent decades in Western countries? *Nutr Metab Cardiovasc Dis*. 2012;22:307–311.
103. Capewell S, Ford ES, Croft JB, et al. Cardiovascular risk factor trends and potential for reducing coronary heart disease mortality in the United States of America. *Bull World Health Organ*. 2010;88:120–130.
104. Bibbins-Domingo K, Grossman DC, Curry SJ, et al. Statin use for the primary prevention of cardiovascular disease in adults: US preventive services task force recommendation statement. *JAMA*. 2016;316:1997–2007.
105. Gosmanova EO, Kovesdy CP. Adherence to antihypertensive medications: is prescribing the right pill enough? *Nephrol Dial Transplant*. 2015;30:1649–1656.
106. Hughes J, Kabir Z, Bennett K, et al. Modelling future coronary heart disease mortality to 2030 in the British Isles. *PLoS One*. 2015;10:e0138044.
107. Scholes S, Bajekal M, Norman P, et al. Quantifying policy options for reducing future coronary heart disease mortality in England: a modelling study. *PLoS One*. 2013;8:e69935.
108. Huffman MD, Lloyd-Jones DM, Ning H, et al. Quantifying options for reducing coronary heart disease mortality by 2020. *Circulation*. 2013;127:2477–2484.
109. Capewell S, O'Flaherty M. Rapid mortality falls after risk-factor changes in populations. *Lancet*. 2011;378:752–753.
110. Tanrikulu MA, Agirbasli M, Berenson G. Primordial prevention of cardiometabolic risk in childhood. *Adv Exp Med Biol*. 2017;956:489–496.
111. Lauer RM, Burns TL, Daniels SR. *Pediatric Prevention of Atherosclerotic Cardiovascular Disease*. New York: Oxford University Press; 2006.
112. Stamler J, Dyer AR, Shekelle RB, et al. Relationship of baseline major risk factors to coronary and all-cause mortality, and to longevity: findings from long-term follow-up of Chicago cohorts. *Cardiology*. 1993;82:191–222.
113. Multiple Risk Factor Intervention Trial Research Group. Multiple Risk Factor Intervention Trial. Risk factor changes and mortality results. *JAMA*. 1982;248:1465–1477.
114. Colquitt JL, Loveman E, O'Malley C, et al. Diet, physical activity, and behavioural interventions for the treatment of overweight or obesity in preschool children up to the age of 6 years. *Cochrane Database Syst Rev*. 2016;3:CD012105.
115. Seery T, Moodie D, Cephus C, et al. Lipid profiles performed at the time of routine physical examinations in 9–11 year olds within Texas Children's Pediatric Associates Primary Care Pediatric Clinics. *J Am Coll Cardiol*. 2014;12(suppl 63):A1461.
116. Din-Dzietham R, Liu Y, Bielo M-V, et al. High blood pressure trends in children and adolescents in national surveys, 1963 to 2002. *Circulation*. 2007;116:1488–1496.
117. Laitinen TT, Pahkala K, Venn A, et al. Childhood lifestyle and clinical determinants of adult ideal cardiovascular health. *Int J Cardiol*. 2013;169:126–132.
118. Dahm CC, Chomistek AK, Jakobsen MU, et al. Adolescent diet quality and cardiovascular disease risk factors and incident cardiovascular disease in middle-aged women. *J Am Heart Assoc*. 2016;5:e003583.
119. Ruys J, Hickie JB. Serum cholesterol and triglyceride levels in Australian adolescent vegetarians. *Br Med J*. 1976;2:87.
120. Yen C-E, Yen C-H, Huang M-C, et al. Dietary intake and nutritional status of vegetarian and omnivorous preschool children and their parents in Taiwan. *Nutr Res*. 2008;28:430–436.
121. Krajčovičová-Kudláčková M, Šimončič R, Béderová A, et al. Influence of vegetarian and mixed nutrition on selected haematological and biochemical parameters in children. *Nahrung*. 1997;41:311–314.
122. Nathan I, Hackett AF, Kirby S. The dietary intake of a group of vegetarian children aged 7–11 years compared with matched omnivores. *Br J Nutr*. 1996;75:533–544.
123. al-Dlaigan YH, Shaw L, Smith AJ. Vegetarian children and dental erosion. *Int J Paediatr Dent*. 2001;11:184–192.
124. Macknin M, Kong T, Weier A, et al. Plant-based no added fat or American Heart Association diets, impact on cardiovascular risk in obese hypercholesterolemic children and their parents. *J Pediatr*. 2015;166:953–959.e3.
125. Melina V, Craig W, Levin S. Position of the academy of nutrition and dietetics: vegetarian diets. *J Acad Nutr Diet*. 2016;116:1970–1980.